

Hubungan Mekanisme koping

by Arif Setyo Upoyo

Submission date: 17-Feb-2023 04:13PM (UTC+0700)

Submission ID: 2016422025

File name: Manuscript_naskah_publikasi_Nirmala_Rumaja_Putri_l1B019036.pdf (1,004.25K)

Word count: 5537

Character count: 29145

Factors Associated with Non Hemorrhagic Post-Stroke Depression

Abstract

Background: Post-stroke patient depression can result in delays in stroke recovery and healing. Several factors can influence post-stroke depression including respondent characteristics, degree of stroke severity, and coping mechanisms. The degree of severity of stroke will cause neurological deficits which can become stressors that will affect individual coping mechanisms. This study aims to analyze the factors associated with non-hemorrhagic post-stroke depression. **Methods:** Analytical observational with a cross-sectional approach. The research was conducted at the Neurology Clinic of RSUD dr. R. Goeteng Taroenadibrata Purbalingga. The population is 435 post-stroke non-hemorrhagic patients. The sample is 104 respondents with the sampling technique using consecutive sampling. The instruments used were the Brief COPE, National Institutes of Health Stroke Scale (NIHSS), and Patient Health Questionnaire-9 (PHQ-9). Bivariate analysis used the Kendall-Tau correlation test. **Results:** The majority of respondents were elderly, male, primary school education level/equivalent, married, live with husband/wife and children, left hemisphere lesion location, had stroke duration >2 years, did not experience recurrent strokes, and had hypertension. The majority of respondents had problem-focused coping, mild stroke severity, and were not depressed. Age, gender, marital status, living together, location of the lesion, duration of stroke, stroke frequency, and comorbidities were not significantly related ($p>0.05$) while education level was significantly associated with post-stroke depression ($p=0.024$). There was no significant correlation between coping mechanisms ($p=0.891$) while stroke severity was associated with post-stroke depression ($p=0.006$). **Conclusions:** There was no relationship between age, gender, marital status, living together, lesion location, stroke duration, frequency of stroke, and comorbidities. There is a relationship between education level and severity of stroke with post-stroke depression.

Keywords: post-stroke depression, stroke severity, coping mechanism, non hemorrhagic, stroke.

INTRODUCTION

Stroke is a cerebrovascular disease which is the second cause of death and the third cause of disability in the world. Stroke occurs when blood vessels that carry oxygen and nutrients to the brain are blocked or ruptured so that brain cells die (American Stroke Association, 2022). According to the World Stroke Organization (2019), there are more than 13.7 million new stroke cases in the world each year. The prevalence of stroke in Indonesia reaches 51.6 per 100,000 people with around 4.3% of cases of disability and about 15-27% of deaths (Tanto et al., 2014).

Stroke causes various physiological and psychological impacts on sufferers. The impact of a stroke on a person's physiology includes loss of movement, thinking, memory, and ability to speak, as well as temporary or permanent loss of sensation (Smeltzer, 2013). Stroke can have a psychological impact in the form of anxiety, changes in self-concept, and depression. The most common psychological impact is depression or which can be called post-stroke depression (Biantoro et al., 2015 in Purba and Utama, 2019).

Globally, post-stroke depression can affect a quarter to a third of stroke patients at any one time (Wubshet et al., 2022). The prevalence of post-stroke depression is around 35-79% depending on the patient selection criteria in the research conducted (Llorcaa et al., 2015). Patients with post-stroke depression generally have more disability, poor rehabilitation outcomes, and increased morbidity and mortality in the first year after stroke onset. Therefore, it is important to know the various factors associated with the occurrence of post-stroke depression (Ibrahimagic et al., 2019).

Factors associated with post-stroke depression include age, gender, neuroticism, family history, level of disability, degree of stroke severity, social support, and history of mental disorders (Shi et al., 2017). According to Reni, Suryani, and Sasmita (2020), factors such as education level, comorbidities, duration of a stroke, and motor disorders are associated with post-stroke depression. In addition, self-management is also related to post-stroke depression (Putri, Upoyo and Purnawan, 2018).

The degree of stroke severity may be one of the most important risk factors for post-stroke depression. The higher the degree of stroke severity, the greater the movement disorders and life barriers which are considered to be negative events in post-stroke patients can increase the occurrence of depression (Shi et al., 2017). However, according to Basit and Mahmudah (2019), the degree of stroke severity is not significantly related to post-stroke depression. In addition, there have been no other studies that have examined the relationship between stroke severity and post-stroke depression in Indonesia.

Post-stroke patients generally experience a loss of body function and health which causes a feeling of helplessness which is usually accompanied by depression. Feelings of helplessness arise based on considering individual responses, coping patterns, and patient characteristics (Nuraliyah and Burmanajaya, 2019). Coping patterns or mechanisms are ways of responding to environmental changes or certain problems (Kozier et al., 2011). According to Carver, Scheier and Weintraub (1989), coping mechanisms are divided into problem-focused coping, emotion-focused coping, and less useful coping. Problem-focused coping is an effort to deal with problems by focusing on solving problems. Emotional-focused coping is emotion-focused coping aimed at minimizing the pressure caused by stressors. Less useful coping is in the form of reducing efforts to overcome problems and diversion to other activities. Based on these coping mechanisms, stroke patients can have various coping mechanisms which are thought to affect post-stroke depression.

There is no research related to the relationship between coping mechanisms and post-stroke depression in Indonesia and there are still differences of opinion regarding the relationship between respondent characteristics and the degree of severity of stroke on post-stroke depression, so the authors investigated the factors associated with non-hemorrhagic post-stroke depression.

METHODS

Design

This research is a quantitative study with a correlational analysis design with a cross-sectional approach.

Population and Sample

The population in this study were non-hemorrhagic post-stroke patients who visited the Neurology Clinic at RSUD dr. R. Goeteng Taroenadibrata Purbalingga in November 2022, namely 435 people. Inclusion criteria in this study included non-hemorrhagic post-stroke patients aged 45-74 years, and willing to be respondents. Exclusion criteria in this study were patients with aphasia, impaired cognitive function, visual impairment, hearing loss, or incomplete data. Sampling in this study uses a consecutive sampling technique. Determining the number of samples was calculated using statistical analysis software G*Power with an effect size of 0.3, a significance level of 0.05, and statistical power of 0.85 which resulted in a total sample in this study of 96 respondents assuming a drop out of 10%, the results the sample calculation is 106 respondents (Faul, Erdfelder, Buchner, & Lang, 2009). After carrying out the data collection process, of the 124 patients encountered, 20 of them did not meet the predetermined inclusion and exclusion criteria with details of 1 person aged <45 years, 12 people aged >74 years, 1 person with aphasia, 1 had impaired cognitive function and there were 5 incomplete data on the lesion location questionnaire. Based on this, the final sample in this study amounted to 104 respondents.

Instruments

The characteristics of the respondents were measured using a questionnaire on the characteristics of the respondents. Coping mechanisms are measured using the COPE Brief compiled by Carver in 1997 based on

the development of Lazarus & Folkman's theory. This COPE brief consists of 14 subscales and each subscale has two items and two questions⁴⁰ so there are 28 question items that have been⁵ adapted from the COPE questionnaire. The COPE brief has been tested for validity by Yussof et al.⁵ (2009) with a Cronbach's alpha value of 0.50-0.80 and a reliability test has been carried out by Arvina⁵ with a Cronbach's alpha value of 0.799 (Arfina, 2017).

The degree of stroke severity was measured using the National Institutes of Health Stroke Scale¹⁸ (NIHSS), which consists of 11 examination items. Assessment can be categorized with a score of <5 (low), a score of 5-14 (moderate), a score of 15-24 (severe), and a score of ≥25 (very severe) (Jojang, Runtuwene and Pertiwi, 2016). The NIHSS instrument is a standard instrument so researchers do not test its validity and reliability.

The instrument used²¹ as a measure of depression is The Patient Health Questionnaire (PHQ-9). This questionnaire was developed by Dr. Kurt Kroenke et. al. From Columbia University in 1999 with a Grant⁵ from Pfizer. This measure describes the level of depression in the last two weeks. The PHQ-9 instrument has been tested for validity and reliability with a Cronbach's alpha value of 0.885.

Collecting Data

The data collection process took place from 28 December 2022-16 January 2023. In this study, researchers were assisted by 6 enumerators. Researchers looked at medical record data⁴⁵ assisted by nurses on duty at the clinic for the initial screening of respondents. including name and age according to predetermined inclusion and exclusion criteria. After that, the researcher gave data on prospective respondents to the enumerator whose job was to collect primary data from these respondents according to the research questionnaire. Next, the researcher and enumerator approached the respondent, introduced themselves, read the informed consent, and asked whether they were willing to become research respondents. Respondents who are willing to be given a consent form become respondents to be signed. Then the researcher read the questionnaire to the respondent to be answered by the respondent. If there are incomplete answers, the researcher will check medical and radiological record data (specifically the location of the lesion).

Ethics

This research has received ethical approval with number 942/EC/KEPK/XII/2022. Research ethics involving humans as respondents has 4 principles that must be applied including respect for the respondent (respect for person), benefit (beneficence), not harming the respondent (non maleficence) and the principle of justice (Masturoh and Temesvari, 2018).

Data Analysis

The data analysis used was the frequency distribution for univariate analysis and the Kendall Tau⁴⁹ correlation test for bivariate analysis. Variables that were used for univariate and bivariate analysis in this study included age, gender, education level, marital status, living together, lesion location, stroke length, stroke frequency, comorbidities, coping mechanisms, stroke severity, and post-stroke depression³. The variables in this study have nominal and ordinal data scales, so the statistical test used is the Kendall Tau Correlation Test with a significance level of 0.05, which is to assess the significance of the relationship between the two variables (Saryono, 2011).

RESULT

Relationship Characteristics of Respondets

The relationship between respondent characteristics and non-hemorrhagic post-stroke depression is presented in the following table.

Table 1. Characteristics of Respondets

Variable	Depression										r	p
	12 Minimal		Mild		Moderate		Moderately severe		Severe			
	n	%	n	%	n	%	n	%	n	%		
Age												
Middle age	20	19,2	12	11,5	3	2,9	0	0,0	0	0,0	0,010	0,910
Elderly	27	26,0	11	10,6	7	6,7	0	0,0	0	0,0		
Young old	14	13,5	7	6,7	1	1,0	1	1,0	1	1,0		
Gender												
Male	34	32,7	10	9,6	8	7,7	1	1,0	1	1,0	0,023	0,807
Female	27	26,0	20	19,2	3	2,9	0	0,0	0	0,0		
Education level												
No school	6	5,8	9	8,7	0	0,0	1	1,0	0	0,0	-	0,024
Elementary school equivalent	24	23,1	13	12,5	7	6,7	0	0,0	0	0,0	0,194	
Middle school equivalent	6	5,8	4	3,8	1	1,0	0	0,0	0	0,0		
High school equivalent	12	11,5	4	3,8	2	1,9	0	0,0	0	0,0		
College	13	12,5	0	0,0	1	1,0	0	0,0	1	1,0		
Marital status												
Married	54	51,9	21	20,2	11	10,6	1	1,0	0	0,0	0,115	0,216
Divorced	2	1,9	1	1,0	0	0,0	0	0,0	0	0,0		
Death divorced	5	4,8	8	7,7	0	0,0	0	0,0	1	1,0		
Living together												
Alone	1	1,0	1	1,0	0	0,0	0	0,0	0	0,0	0,026	0,763
Child	6	5,8	6	5,8	0	0,0	0	0,0	1	1,0		
Husband and wife	9	8,7	0	0,0	2	1,9	0	0,0	0	0,0		
Husband/Wife and Children	30	28,8	17	16,3	5	4,8	0	0,0	0	0,0		
Other	15	14,4	6	5,8	4	3,8	1	1,0	0	0,0		
Lesion Location												
Right Hemisphere	27	26,0	13	12,5	6	5,8	1	1,0	1	1,0	-	0,330
Left Hemisphere	31	29,8	17	16,3	5	4,8	0	0,0	0	0,0	0,091	
Right-Left Hemisphere	3	2,9	0	0,0	0	0,0	0	0,0	0	0,0		
Duration of Stroke												
<1 year	18	17,3	15	14,4	4	3,8	0	0,0	0	0,0	-	0,245
1-2 years	7	6,7	4	3,8	1	1,0	0	0,0	0	0,0	0,106	
>2 years	36	34,6	11	10,6	6	5,8	1	1,0	1	1,0		
Stroke Frequency												
First Stroke	48	46,2	22	21,7	9	8,7	0	0,0	0	0,0	0,085	0,371
Recurrent Strokes	13	12,5	8	7,7	2	1,9	1	1,0	1	1,0		
Comorbidities												
Nothing	13	12,5	3	2,9	3	2,9	0	0,0	0	0,0	0,075	0,392
Diabetes mellitus	2	1,9	2	1,9	0	0,0	0	0,0	0	0,0		
Hypertension	25	24,0	16	15,4	2	1,9	0	0,0	0	0,0		
Heart disease	0	0,0	0	0,0	2	1,9	0	0,0	0	0,0		
Other	21	20,2	9	8,7	4	3,8	1	1,0	1	1,0		

Based on table 1, it is known that of the 104 respondents, the majority were elderly (43.3%) and male (51.9%). The majority of respondents have the last education level elementary school equivalent (42.3%), are married (83.7%), and live with their husband/wife and their children (50.0%). Most of the respondents had a stroke lesion location in the left hemisphere (51.0%) with a stroke duration of more than two years (52.9%) and a first attack stroke (76.0%). The majority of respondents in this study had comorbidities, namely hypertension (41.3%).

The results of bivariate analysis on the variables age, sex, marital status, living together, lesion location, stroke duration, stroke frequency, and comorbidities obtained $p > 0.05$ indicating that there was no significant relationship between these variables and depression. While the variable level of education has a value of $p = 0.024$ which indicates a significant relationship with post-stroke depression. The value of r or correlation coefficient of education level with depression in stroke patients is -0.194. This shows that there is a negative direction of the relationship and the strength of the relationship is "very weak".

Correlation between Coping Mechanisms and Stroke Severity Degree with Non-Hemorrhagic Post-Stroke Depression

The relationship between coping mechanisms and stroke severity with non-hemorrhagic post-stroke depression is described in table 2.

Table 2. Coping Mechanisms and Stroke Severity Degree

Variable	Depresi										<i>r</i>	<i>p</i>
	Minimal		Mild		Moderate		Moderately severe		Severe			
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%		
Coping mechanism												
Problem Focused Coping (PFC)	34	32,7	16	15,4	6	5,8	0	0,0	1	1,0	0,013	0,891
Emotion Focused Coping (EFC)	26	25,0	14	13,5	5	4,8	1	1,0	0	0,0		
PFC=EFC	1	1,0	0	0,0	0	0,0	0	0,0	0	0,0		
Stroke severity degree												
Mild	54	51,9	25	24,0	5	4,8	1	1,0	0	0,0	0,260	0,006
Moderately severe	7	6,7	5	4,8	6	5,8	0	0,0	1	1,0		

Based on table 2 it is known that the majority of respondents have a coping mechanism, namely problem focused coping as many as 56 respondents (54.8%). Most of the respondents had mild stroke severity, 85 respondents (81.7%) and the majority of respondents had minimal or no depression, 61 respondents (58.7%). The results of the Kendall tau test on coping mechanism variables have a value of $p = 0.891$ ($p < 0.05$) which means there is no significant relationship between coping mechanism variables and post-stroke depression. The stroke severity variable has a value of $p = 0.006$ ($p < 0.05$) which indicates a significant relationship between stroke severity and depression in post-stroke patients. The value of r or the correlation coefficient of the severity of stroke with depression in stroke patients is 0.260. This shows that there is a positive relationship direction with a "sufficient" relationship strength.

DISCUSSIONS

The results of the analysis of the relationship between the characteristics of respondents with non-hemorrhagic post-stroke depression showed mixed results. Age is not associated with post-stroke depression. This is possible due to other factors such as living with family. Depression in the elderly is caused by the loneliness felt by the elderly who live alone, without children, low health conditions, low education levels and low self-confidence (Herawati & Deharnita, 2019). In line with the research of Khedr et

al. (2020), age does not show a significant relationship with post-stroke depression because there are other factors such as family support and coping mechanisms.

Gender was not associated with post-stroke depression. Gender is not a risk factor for post-stroke depression. Post-stroke depression occurs more frequently in women, whereas in this study the majority of respondents were male (Khedr et al., 2020). This might be one of the reasons for the absence of a relationship between gender and post-stroke depression in this study.

Respondents' level of education is related to post-stroke depression with a negative direction and weak relationship strength. This shows that the higher the level of education of the respondents, the lower the level of depression. Someone with a high level of education will find it easier to overcome depression. This is because education is the main provision for cognitive development which will become a mediator when facing a certain incident (Setiawati and Ismahmudi, 2020). Education can influence a person's behavior regarding lifestyle, motivation, and attitudes that play a role in building health. The higher a person's education, the easier it is to receive information and the more knowledge one has (Herawati and Dehamita, 2019). Someone with minimal knowledge will hinder the development of one's attitude towards acceptance, information, and values so that it is prone to depression (Nilamsari and Handayani, 2014). In line with the research of Khedr et al. (2020) that the majority of respondents who experience depression have a low educational level.

Marital status and living together were not associated with post-stroke depression. The majority of respondents are married and live with their families, so it is possible to get family information support. Family information support influences the healing and recovery of post-stroke patients (Kurniawan and Wibowo, 2020). In line with the results of the study (Purba and Utama, 2019) which showed that there was no significant relationship between marital status and post-stroke depression.

The location of the lesion has no significant relationship with post-stroke depression. In this study, it was also found that respondents had lesions located on the right and left hemispheres. Although some suggest that the location of the lesion in the left hemisphere is associated with post-stroke depression, this is still being debated. This is associated with emotion and language functions dominated by the left hemisphere of the brain. The parts of the brain that regulate emotions, namely the frontal lobe and basal ganglia, tend to experience changes that lead to depressive symptoms (Shi et al., 2017). In line with research (Pongsilurang et al., 2020) which states that there is no relationship between the location of stroke lesions and depression.

Stroke duration and stroke frequency did not have a significant relationship with post-stroke depression. According to Hayulita and Sari (2014) long-suffering strokes and the frequency of strokes will make patients more hopeless about their illness, feel helpless even though they have different defense mechanisms, they will still find it difficult to deal with the stressor. The majority of respondents had coping mechanisms as a defense that varied and the majority were not depressed while the majority had suffered a stroke for more than 2 years. This is the reason for the absence of a significant relationship between the two variables in this study.

Comorbidities are not associated with post-stroke depression. According to Hayulita (2014), co-morbidities can be the etiology or complication of stroke patients, but the coping mechanisms for each patient are different, causing depression.

Coping mechanisms are not associated with post-stroke depression. Coping mechanisms will play a role in dealing with the stressors caused by the stroke. Everyone has their own coping mechanism and will be adapted to their individual circumstances (Loupatty, Ranimpi and Rayanti, 2019). Rutter (1981) states that the most effective coping of stress is according to the type of stress and situation. Thus, the coping mechanism owned by the respondent is not absolute in just one type of coping mechanism and can change at any time depending on the stressor being faced. Research conducted by Supriyati, Noerhidajati and Julianti (2020) explains that problem focused coping mechanisms tend to be used if respondents think problems can be controlled so that to deal with stressors that cause depression they try to change or overcome the problems they face. This research was conducted in a clinic and patients who visited had the goal of curing their illness. This is one of the signs that patients use problem focused coping. Emotion focused coping mechanisms are used by respondents who perceive a problem as difficult to control, so that in dealing with stressors that cause depression, they regulate their emotional responses to adjust to stressful situations. This type of coping may be used more by patients who do not visit treatment or rehabilitation because they feel they can adapt to their condition. In this study, the coping mechanisms owned by the respondents were not categorized into adaptive and maladaptive, so that all of the coping mechanisms they had could play a role in dealing with the stressors they encountered.

The degree of severity of stroke with post-stroke depression shows a significant relationship with the direction of a positive relationship and the strength of the relationship is sufficient. This shows that the higher the degree of stroke severity, the higher the level of depression. The degree of stroke severity indicates how severe the neurologic deficit is experienced by the patient. The higher the neurological deficit, the more disorders the patient will experience which can then affect the patient's quality of life (Maharani, July, and Hermawan, 2021). The degree of stroke severity affects the level of disability and independence, which are factors that need to be considered with the occurrence of post-stroke depression (Shi et al., 2017). Impaired motor function in the form of limb weakness can lead to an inability to self-care and reduce mobility which can hinder the fulfillment of daily life activities. Thus, in meeting their basic needs, stroke patients need help from a caregiver (Sonatha, 2012). Stroke patients who have a low level of independence will have a significant relationship with emotional distress. One sign of emotional distress that often occurs in stroke patients is depression (Thomas and Lincoln, 2008).

Dependence in stroke patients is also associated with patient self-esteem. The higher the level of dependency caused by the high degree of stroke severity, the higher the patient's self-esteem, and vice versa (Fadlulloh, Upoyo, and Hartanto, 2014). Physical limitations, reduced activities of daily living, and inability to work arise because of a stroke which affects a person and makes him sad (Suharsono, 2022). Stroke patients rate themselves as low, feel tired, are physically disabled, and have a low quality of life that will support the occurrence of post-stroke depression (Sibbritt et al., 2022). The prevalence of post-stroke depression is higher in patients with reduced mobility, high functional dependence, and more severe functional disabilities (Kumar et al., 2020). In this study, the majority of post-stroke depression was either minimally depressed or not depressed. This is because the degree of severity of stroke in the respondents in this study was the majority of the degree of severity of mild stroke.

CONCLUSIONS & NURSING IMPLICATION

Age, gender, marital status, living together, location of the lesion, duration of a stroke, stroke frequency, and morbidities did not show a significant relationship with post-stroke depression, while education level had a significant relationship with post-stroke depression. Coping mechanisms are not associated with post-stroke depression while the degree of stroke severity is associated with post-stroke depression. Based on this study, it can be seen that the level of education and stroke severity are associated with non-hemorrhagic post-stroke depression. Nurses are expected to provide health education related to how to prevent or reduce depression to increase knowledge due to low education levels in post-stroke patients. In addition, nurses can also motivate patients to always take medication regularly so that they can reduce the severity of a stroke and prevent or reduce the occurrence of depression after non-hemorrhagic stroke.

ACKNOWLEDGEMENT

In the process of this research, the authors received assistance from various parties involved. Therefore, authors would like to thank the Dean of the Faculty of Health Sciences and all Lecturers in the Department of Nursing, Jenderal Soedirman University, Director of RSUD dr. R. Goeteng Taroenadibrata Purbalingga and his staff and staff, all enumerators, respondents, and other parties that cannot be mentioned one by one.

REFERENCES

- American Stroke Association. (2022). About Stroke | American Stroke Association. *American Stroke Association*.
- Amila, Sinaga, J. & Sembiring, E. 2018, "Pencegahan Stroke Berulang Melalui Pemberdayaan Keluarga dan Modifikasi Gaya Hidup," *ABDIMAS*, vol. 22, no. 2. <https://doi.org/10.31294/jabdimas.v2i2.6120>
- Arfina, A. (2017). *Hubungan Mekanisme Koping dan Disabilitas dengan Keputusan pada Pasien Stroke di Rumah Sakit Kota Medan*. Universitas Sumatra Utara.
- Basit, M., & Mahmudah, R. (2019). Analisis Faktor Kejadian Depresi Pada Klien Pasca Stroke Suku Banjar Banjarmasin. *Dinamika Kesehatan Jurnal Kebidanan Dan Keperawatan*, 10(2), 730–744. <https://doi.org/10.33859/dksm.v10i2.508>
- Carver, C. S., Scheier, M. F., & Weintraub, J. K. (1989). Assessing Coping Strategies: A Theoretically Based Approach. *Journal of Personality and Social Psychology*, 56(2), 267–283. <https://doi.org/10.1037/0022-3514.56.2.267>
- Fadlulloh, S. F., Upoyo, A. S., & Hartanto, Y. D. (2014). Hubungan Tingkat Ketergantungan Dalam Pemenuhan Aktivitas Kehidupan Sehari-Hari (AKS) Dengan Harga Diri Penderita Stroke. *Jurnal Keperawatan Soedirman*, 9(2), 134–145. <http://dx.doi.org/10.20884/1.jks.2014.9.2.579>
- Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses using G*Power 3.1: Tests for correlation and regression analyses. *Behavior Research Methods*, 41(4), 1149–1160. <https://doi.org/10.3758/BRM.41.4.1149>
- Hayulita, S., & Sari, D. R. (2014). Faktor-Faktor yang Berhubungan Dengan Depresi Pada Pasien Paska Stroke di Ruang Rawat Jalan Rumah Sakit Stroke Nasional (RSSN) Bukittinggi Tahun 2014. *Yasri Sumbar*, (April). Diambil dari <http://ejournal.stikesyarsi.ac.id/index.php/JAV1N1/article/download/32/123>
- Herawati, N., & Deharnita. (2019). Hubungan Karakteristik dengan Kejadian Depresi pada Lansia. *Jurnal Keperawatan Jiwa*, 7(2), 183–190. <https://doi.org/10.26714/jkj.7.2.2019.185-192>
- Ibrahimagic, O. C., Smajlovic, D., Kunic, S., Dostovic, Z., Custovic, A., Sehanovic, A., & Kojic, B. (2019). Post-

- Stroke Depression. *Mater Sociomed*, 31(1), 31–41. <https://doi.org/10.5455/msm.2019.31.31-34>
- Jojang, H., Runtuwene, T., & Pertiwi, J. M. (2016). Perbandingan NIHSS pada Pasien Stroke Hemoragik dan Non-hemoragik yang Rawat Inap di Bagian Neurologi RSUP Prof. Dr. R. D. Kandou Manado. *Jurnal e-Clinic*, 4(1). <https://doi.org/10.35790/ed.4.1.2016.12111>
- Khedr, E. M., Abdelrahman, A. A., Desoky, T., Zaki, A. F., & Gamea, A. (2020). Post-Stroke Depression: Frequency, Risk Factors, and Impact on Quality Life Among 103 Stroke Patients- Hospital-Based Study. *The Egyptian Journal of Neurology, Psychiatry and Neurosurgery*, 56(66). <https://doi.org/10.1186/s41983-020-00199-8>
- Kozier, B., Erb, G., Berman, A., & Snyder, S. J. (2011). *Buku Ajar Fundamental Keperawatan : Konsep, Proses, & Praktik* (7 ed.; D. Widiarti, B. Subekti, & A. O. Tampubolon, Ed.). Jakarta: EGC.
- Kumar, R., Kataria, N., Kumar, N., Kumar, M., & Bahurupi, Y. (2020). Poststroke depression among stroke survivors in Sub-Himalayan region. *Journal of Family Medicine and Primary Care*, 9(7), 3508–3513. https://doi.org/10.4103/jfmpc.jfmpc_444_20
- Kurniawan, M. B., & Wibowo, T. A. (2020). Hubungan antara Dukungan Informasi Keluarga dengan Depresi pada Pasien Pasca Stroke di Puskesmas Remaja Samarinda. *Borneo Student Research*, 1(2).
- Llorca, G. E., Castilla-Guerrab, L., Morenoc, M. C. F., Dobladoa, S. R., & Hernández, M. D. J. (2015). Post-stroke depression: an update. *Neurologia*, 30(1), 23–31. <https://doi.org/10.1016/j.nrleng.2012.06.006>
- Loupaty, S. N., Ranimpi, Y. Y., & Rayanti, R. E. (2019). Respon Psikososial dan Strategi Koping Pasien Stroke dalam Konteks Budaya Ambon. *Jurnal Kesehatan*, 10(3), 480–488. <https://doi.org/10.26630/jk.v10i3.1121>
- Maharani, T., Juli, C., & Hermawan, A. N. (2021). Karakteristik Luaran Klinis Neurologis Pasien Stroke Iskemik berdasarkan NIHSS. *Jurnal Kedokteran Syiah Kuala*, 21(3), 212–222. <https://doi.org/10.24815/jks.v21i3.20578>
- Masturoh, I., & Temesvari, N. A. (2018). *Metodologi Penelitian Kesehatan*. Jakarta: Kemenkes RI. <https://doi.org/10.24815/jks.v21i3.20646>
- Nilamsari, & Handayani, N. (2014). Tingkat Pengetahuan Akan Memengaruhi Tingkat Depresi Penderita Kanker. *Jurnal Ilmiah Kesehatan*, 7(2), 107–113. <https://doi.org/10.33086/jhs.v7i2.498>
- Nuraliyah, S., & Burmanajaya, B. (2019). Mekanisme Koping dan Respon Ketidakberdayaan pada Pasien Stroke. *Jurnal Riset Kesehatan*, 11(1), 38–43.
- Pongsilurang, J. M., Pieter, H., Wungouw, L., Setiono, K. W., & Kupang, J. (2020). Hubungan Letak Lesi Stroke Dengan Kejadian Depresi Pasca Stroke Pada Pasien Stroke Di Rsud Prof Dr W Z Johannes Kupang. *Cendana Medical Journal*, 19(April), 51–57. Diambil dari https://r.search.yahoo.com/_ylt=Awrwv7Nh.Vh7DQA7QDLQwx.;_ylu=Y29sbwNzZzMEcG9zAzEEdnRpZAMEc2VjA3Ny/RV=2/RE=1642461261/RO=10/RU=https%3A%2F%2Fjurnal.undana.ac.id%2FCMJ%2Fissue%2Fview%2F286/RK=2/RS=mQiwYondoQeso5omjXThDiPK.0Y-
- Purba, M. M., & Utama, N. R. (2019). Disabilitas Klien Pasca Stroke terhadap Depresi. *Jurnal Kesehatan*, 10(3), 346. <https://doi.org/10.26630/jk.v10i3.1539>
- Putri, C. K., Upoyo, A. S., & Purnawan, I. (2018). *Hubungan Self Management dengan Depresi pada Pasien Pasca Stroke di Poliklinik Syaraf RSUD Dr. R. Goeteng Taroenadibrata Purbalingga*. Universitas Jenderal Soediman.
- Reni, Suryani, & Sasmita, H. (2020). Analisa Faktor-Faktor yang Berhubungan dengan Tingkat Depresi Pasca Stroke di Rumah Sakit Stroke Nasional (RSSN) Bukittinggi. *Jurnal Kesehatan Mercusuar*, 3(2), 24–31. <https://doi.org/https://doi.org/10.36984/jkm.v3i2.84>

- Rutter, M. (1981). Stress, coping, and development: some issues and some questions. *Child Psychology & Psychiatry & Allied Disciplines*, 22(4), 323–356. <https://doi.org/https://psycnet.apa.org/doi/10.1111/j.1469-7610.1981.tb00560.x>
- Saryono. (2011). *Metodologi Penelitian*. Purwokerto: UPT. Percetakan dan Penerbitan UNSOED.
- Setiawati, T. I., & Ismahmudi, R. (2020). Hubungan Pekerjaan dan Pendidikan dengan Tingkat Depresi pada Lansia di Posyandu Lansia Wilayah Kerja Puskesmas Wonorejo Samarinda. *Borneo Student Research*, 1(3).
- Shi, Y., Yang, D., Zeng, Y., & Wu, W. (2017). Risk Factors for Post-stroke Depression: A Meta-analysis. *Front Aging Neurosci*, 9(2018). <https://doi.org/10.3389/fnagi.2017.00218>
- Sibbritt, David, Bayes, J., Peng, W., Maguire, J., & Adams, J. (2022). Associations Between Fatigue and Disability, Depression, Health-Related Hardiness and Quality of Life in People with Stroke. *Journal of Stroke and Cerebrovascular Disease*, 31(7). <https://doi.org/10.1016/j.jstrokecerebrovasdis.2022.106543>
- Smeltzer, S. C. (2013). *Keperawatan Medikal Bedah: Brunner & Suddarth; Alih Bahasa, Devi Yulianti, Amelia Kimin* (12 ed.; E. A. Mardela, Ed.). Jakarta: EGC.
- Sonatha, B. (2012). *Hubungan tingkat pengetahuan dengan sikap keluarga dalam pemberian perawatan pasien pasca stroke*. Universitas Indonesia.
- Suharsono. (2022). Hubungan Antara Harga Diri dengan Tingkat Depresi pada Pasien Pasca Stroke. *Jurnal Kesehatan Karya Husada*, 10(2).
- Supriyati, R., Noerhidajati, E., & Julianti, H. P. (2020). Hubungan Antara Mekanisme Koping Dengan Derajat Depresi Pada Caregiver Pasien Skizofrenia Studi di Unit Rawat Jalan Rumah Sakit Jiwa Amino Gondo Hutomo Provinsi Jawa Tengah. *Medica Hospitalia*, 7(2). <https://doi.org/https://doi.org/10.36408/mhjcm.v7i2.520>
- Tanto, C., Liwang, F., Hanifati, S., & Pradipta, E. A. (2014). *Kapita Selekta Kedokteran* (4 ed.). Jakarta: Media Aesculapius.
- Thomas, S. A., & Lincoln, N. B. (2008). Predictors of Emotional Distress After Stroke. *Stroke*, 39(4), 1240–1245. <https://doi.org/10.1161/STROKEAHA.107.498279>
- Towfighi, A., Ovbiagele, B., El Hussein, N., Hackett, M. L., Jorge, R. E., Kissela, B. M., ... Williams, L. S. (2017). Poststroke Depression: A Scientific Statement for Healthcare Professionals from the American Heart Association/American Stroke Association. *Stroke*. <https://doi.org/10.1161/STR.0000000000000113>
- World Stroke Organization. (2019). Global Stroke Fact Sheet. In *World Stroke Organization*.
- Wubshet, T. Y., Geberemichael, S. G., Adilo, T. M., Arusi, T. T., Gutulo, M. G., Assefa, D. Z., & Asfaw, M. W. (2022). Prevalence and Associated Factors of Poststroke Depression among Outpatient Stroke Patients Who Have a Follow-Up at the Outpatient Neurology Clinic of Zewditu Memorial Hospital in Addis Ababa, Ethiopia. *Depression Research and Treatment*, 2022. <https://doi.org/10.1155/2022/9750035>

Hubungan Mekanisme koping

ORIGINALITY REPORT

22%

SIMILARITY INDEX

18%

INTERNET SOURCES

15%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1

"1st Annual Conference of Midwifery", Walter de Gruyter GmbH, 2020

Publication

3%

2

eprints.nottingham.ac.uk

Internet Source

2%

3

epubs.surrey.ac.uk

Internet Source

2%

4

www.ncbi.nlm.nih.gov

Internet Source

1%

5

garuda.kemdikbud.go.id

Internet Source

1%

6

jurnal.mercubaktijaya.ac.id

Internet Source

1%

7

www.science.gov

Internet Source

1%

8

scholarworks.waldenu.edu

Internet Source

1%

9

Submitted to Nightingale College - School of Nursing

1%

10	www.medrxiv.org Internet Source	1 %
11	Submitted to University of Portsmouth Student Paper	<1 %
12	bmcpsy psychiatry.biomedcentral.com Internet Source	<1 %
13	theses.gla.ac.uk Internet Source	<1 %
14	medicahospitalia.rskariadi.co.id Internet Source	<1 %
15	proceeding.tenjic.org Internet Source	<1 %
16	"Neurobiological and Psychological Aspects of Brain Recovery", Springer Science and Business Media LLC, 2017 Publication	<1 %
17	Amanda Candraning Pratiwi, Haryatiningsih Purwandari, Meivita Dewi Purnamasari. "The effects of the early oral feeding on preterm infant's length of hospital stay", AIP Publishing, 2023 Publication	<1 %
18	Submitted to Ivy Tech Community College Central Office Student Paper	<1 %

19

Submitted to Buckinghamshire Chilterns
University College

Student Paper

<1 %

20

medica-musc.researchcommons.org

Internet Source

<1 %

21

Submitted to Adtalem Global Education, Inc.

Student Paper

<1 %

22

jurnal.payungnegeri.ac.id

Internet Source

<1 %

23

archive.org

Internet Source

<1 %

24

jfmipc.com

Internet Source

<1 %

25

journal.unnes.ac.id

Internet Source

<1 %

26

lucris.lub.lu.se

Internet Source

<1 %

27

Submitted to Laureate Higher Education
Group

Student Paper

<1 %

28

Mei - Yung Leung, Anita M. M. Liu, Maggie
Mei - ki Wong. "Impact of stress - coping
behaviour on estimation performance",
Construction Management and Economics,
2006

Publication

<1 %

29	newinera.com Internet Source	<1 %
30	Submitted to University of Edinburgh Student Paper	<1 %
31	ojs.stikespanritahusada.ac.id Internet Source	<1 %
32	Andrew R. Newberg, Dmitry S. Davydow, Hochang B. Lee. "Cerebrovascular disease basis of depression: Post-stroke depression and vascular depression", International Review of Psychiatry, 2009 Publication	<1 %
33	Jufang Li, Yun Li, Ping Li, Maiyun Ye. "Early Symptom Measurement of Post-Stroke Depression: Development and validation of a new short version", Journal of Advanced Nursing, 2019 Publication	<1 %
34	Rebecca El-Helou, Jeffrey M. Rogers, Brooke Ryan, Dianne L. Marsden, Ann Winkler, Andrew Baillie, Ian Kneebone. "Evaluating the feasibility and acceptability of the Mood Assessment Post-Stroke (MAPS) mood screening training", Brain Impairment, 2022 Publication	<1 %
35	Submitted to Royal College of Surgeons in Ireland	<1 %

-
- | | | |
|----|---|------|
| 36 | journal.ppnijateng.org
Internet Source | <1 % |
|----|---|------|
-
- | | | |
|----|---|------|
| 37 | jurnal.stikes-alinsyirah.ac.id
Internet Source | <1 % |
|----|---|------|
-
- | | | |
|----|---|------|
| 38 | link.springer.com
Internet Source | <1 % |
|----|---|------|
-
- | | | |
|----|---|------|
| 39 | ojs.dinamikakesehatan.unism.ac.id
Internet Source | <1 % |
|----|---|------|
-
- | | | |
|----|---|------|
| 40 | Priyo Purnomo As'hab, Budi Anna Keliat, Ice Yulia Wardani. "The Effects of Acceptance and Commitment Therapy on Psychosocial Impact and Adherence of Multidrug-Resistant Tuberculosis Patients", Journal of Public Health Research, 2022
Publication | <1 % |
|----|---|------|
-
- | | | |
|----|---|------|
| 41 | amedeo.com
Internet Source | <1 % |
|----|---|------|
-
- | | | |
|----|---|------|
| 42 | hdl.handle.net
Internet Source | <1 % |
|----|---|------|
-
- | | | |
|----|---|------|
| 43 | ijamscr.com
Internet Source | <1 % |
|----|---|------|
-
- | | | |
|----|---|------|
| 44 | journal.um-surabaya.ac.id
Internet Source | <1 % |
|----|---|------|
-
- | | | |
|----|--|--|
| 45 | jurnal.htp.ac.id | |
|----|--|--|

<1 %

46

jurnal.unsyiah.ac.id

Internet Source

<1 %

47

koreascience.or.kr

Internet Source

<1 %

48

vdoc.pub

Internet Source

<1 %

49

www.healio.com

Internet Source

<1 %

50

Juliana Gracia G.E.P Massie, Ratna Sitorus, I Made Kariasa, Yunisar Gultom, Maya Khairani, Noor Wulandari. "RISK OF POST STROKE PNEUMONIA DURING HOSPITALIZATION", Jurnal Mitra Kesehatan, 2021

Publication

<1 %

51

Haiyan Liu, Juncal Pu, Qinxiang Zhou, Lining Yang, Dingqun Bai. "Peripheral blood and urine metabolites and biological functions in post-stroke depression", Metabolic Brain Disease, 2022

Publication

<1 %

52

Shrikant D Pande, May Thiri Lwin, Kaung Myat Kyaw, Aye Aye Khine, Aye Aye Thant, May Myat Win, Julie Morris. "Post-stroke Seizure;

<1 %

do the locations, types and managements of stroke matter?", Epilepsia Open, 2018

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On