

Publikasi Puteri Aulia Azzahra.docx

by cek turnitin

Submission date: 02-Jan-2024 01:41AM (UTC-0600)

Submission ID: 2264823301

File name: Publikasi_Puteri_Aulia_Azzahra.docx (91.93K)

Word count: 2673

Character count: 15816

Correlation Between Blood Pressure at Emergency Room and Mortality in Acute Non-Hemorrhagic Stroke

Puteri Aulia Azzahra¹, Tri Wahyuliati^{2,*}

¹Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Special Region of Yogyakarta, Indonesia

² Department of Neurology, Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta, Special Region of Yogyakarta, Indonesia

DATA OF ARTICLE:

Received:

Reviewed:

Revised:

Accepted: ...

*CORRESPONDENCE:

Institutional@author

DOI:

.....

TYPE OF ARTICLE:

Research

Abstract: According to data from the World Stroke Organization (WSO), the annual incidence of stroke is around 13.7 million new cases, with an estimated 5.5 million deaths among those impacted. Stroke caused by complications of hypertension causes 51% of the 9.4 million deaths worldwide. The neurological severity of stroke patients is associated with a decrease in systolic blood pressure of 10 mmHg for every value below 150 mmHg. The purpose of this study is to examine the correlation between blood pressure levels and mortality in acute non-hemorrhagic stroke patients. This study uses cross-sectional research methods to conduct observational research. The study included a total of 60 patients with acute non-hemorrhagic strokes who were admitted to PKU Muhammadiyah Gamping Hospital. The patients were all over the age of 50 and met the specified inclusion and exclusion criteria. Data analysis using univariate tests and Spearman's bivariate tests. The results showed that there is no significant correlation between blood pressure at hospital admission and mortality in non-hemorrhagic stroke patients, as evidenced by a p-value of 0.5. The conclusion is that there is no meaningful correlation between blood pressure on admission to the hospital and mortality in patients with acute non-hemorrhagic strokes.

Keywords: Stroke, blood pressure, mortality

INTRODUCTION

According to data from the World Stroke Organization (WSO), the annual incidence of stroke is around 13.7 million new cases, with an estimated 5.5 million deaths among those impacted. Stroke is the primary factor leading to death across all age groups, accounting for 15.4% of all deaths.¹ A non-hemorrhagic stroke, also known as an ischemic stroke, refers to the necrosis of brain tissue resulting from the interruption of blood supply to the brain. This interruption can be caused by the blockage of cerebral or cervical arteries, or less commonly, by the blockage of cerebral veins.² Thrombosis, embolism, and reduced perfusion pressure are the three main causes of ischemic stroke. Thrombosis is the condition when blood flow is decreased or completely stopped because of the obstruction of blood vessels in a specific area. The vascular disorders that contribute to thrombosis are arteritis, atherosclerosis, vascular dissection, bleeding throughout atherosclerotic plaque, and fibromuscular dysplasia. Atherosclerosis is a major vascular condition.³

Blood pressure is acquired through the mechanism of the heart pumping blood. Systolic blood pressure is the result of the contraction of the ventricles, forcing blood into the arteries. Diastolic blood pressure refers to the pressure exerted on the arterial walls when the ventricles of the heart are in a relaxed state and are being filled with blood from the atria. The average blood pressure of healthy young individuals is 120/80 mmHg.⁴ Hypertension, often known as high blood pressure, is the primary risk factor associated with stroke.⁵ Hypertension can cause disruption of blood circulation in the brain due to thickening and remodeling of blood vessels, which will result in a reduction in the diameter of blood vessels and trigger atherosclerosis.⁶ Orthostatic hypotension can cause reduced venous return, so cardiac output

will also decrease. Decreased cardiac output can result in blood pooling in the lower body and reducing blood flow to the brain.⁷

Stroke patients with elevated blood pressure are more likely to exhibit severe neurological deficits in comparison to stroke patients with lower blood pressure.⁸ Neurological severity and clinical consequences, including earlier death, are connected with a reduction in systolic blood pressure of 10 mmHg below 150 mmHg.⁹ The purpose of this study is to examine the correlation between blood pressure levels and mortality in acute non-hemorrhagic stroke patients.

MATERIALS AND METHOD

This study uses cross-sectional research methods to conduct observational research. The independent variable used in the study was the blood pressure recorded at the beginning of hospital admission, while the dependent variable was the mortality incidence among patients with acute non-hemorrhagic stroke. Blood pressure is divided into 5 categories: low (<90/0 mmHg), normal (<120/80 mmHg), elevated (120-129/80 mmHg), hypertension stage I (130-139/80-89 mmHg), and hypertension stage II (>140/90 mmHg).¹⁰ The research subjects obtained in this study were 60 medical records of patients who were admitted to the emergency room at PKU Muhammadiyah Gamping Hospital. The subjects involved were selected based on inclusion criteria, including non-hemorrhagic stroke patients aged >50 years, and the exclusion criteria were a history of previous strokes. Data analysis using SPSS with univariate tests and Spearman's bivariate tests. This research was conducted at PKU Muhammadiyah Gamping Hospital from August 2023 to September 2023.

RESULT

The data is categorized based on characteristics using the SPSS bivariate descriptive test to determine frequencies and percentages, as shown in Table 1.

Table 1 Subject Characteristics

Characteristics	Percentage			
	Died		Not died	
	n	%	n	%
Gender				
Male	3	5	27	45
Female	7	11,7	23	38,3
Age				
51-60 years old	2	3,3	15	25
61-70 years old	2	3,3	16	26,7
71-80 years old	1	1,7	11	18,3
81-90 years old	3	5	6	10
91 years old and above	2	3,3	2	3,3
Blood pressure				
Low	0	0	2	3,3
Normal	1	1,7	1	1,7
Elevated	0	0	1	1,7
Hypertension stage I	2	3,3	6	10
Hypertension stage II	7	11,7	40	66,7

The study included a total of 60 subjects who had non-hemorrhagic strokes. The participants had an equal distribution according to gender, with a 50% representation of males and a 50% representation of females. The majority of the participants, representing 30% of the total, were in the age range of 61 to 70 years old. Most of the subjects came under the classification of hypertension stage II (78,3%).

The correlation between blood pressure and mortality

The correlation between blood pressure and mortality was analyzed through the Spearman's rank correlation test on SPSS. This test was used to determine the correlation between different blood pressure categories, as presented in Table 2.

Table 2 The correlation between blood pressure and mortality

Table 2 The correlation between blood pressure and mortality							
Blood pressure	Percentage				Total	p	r
	Died		Not died				
	n	%	n	%			
Low	0	0	2	3,3	2 (3,3%)	0,521	0,084
Normal	1	1,7	1	1,7	2 (3,3%)		
Elevated	0	0	1	1,7	1 (1,7%)		
Hypertension stage I	2	3,3	6	10	8 (13,3%)		
Hypertension stage II	7	11,7	40	66,7	47 (78,3%)		
Total	10	16,7	50	83,3	60 (100%)		

The results of the correlation test between blood pressure and mortality using the Spearman test obtained an r value of 0.084 with a p value of 0.521. The p value is > 0.05 , so there is no correlation between blood pressure at hospital admission and mortality in acute non-hemorrhagic stroke patients, and the relationship is very weak.

DISCUSSION

The characteristic distribution results of the study indicate that women have a higher incidence of mortality among non-hemorrhagic stroke patients. This is consistent with previous studies that showed higher mortality rates in women with ischemic strokes. On average, female stroke patients have a higher risk of death than men.¹¹ The subjects of this study are over 50 years of age and therefore belong to the elderly category. The incidence of stroke in women increased in correlation with increasing age and aligned with the beginning of menopause and a decrease in levels of female sex hormones. Estrogen helps prevent atherosclerotic plaques from forming in the blood vessels of the brain. Estrogen production decreases before and during menopause.¹²

Mortality in ischemic stroke patients ranges from 51 years old to over 91 years old. This is consistent with previous research showing that people aged ≥ 55 years are 3,23 times more likely to have a stroke. The risk of stroke tends to increase with age, and the likelihood of having the disease doubles after age 55.¹³ The age of a patient is a significant determinant of the outcome or prognosis of an ischemic stroke. The process of aging results in the progressive aging of blood vessels, which can cause the endothelium to thicken as a result of atherosclerosis.⁶ Decline in vascular function occurs in older people. Vascular elasticity is reduced, especially in the intima where endothelial thickening occurs, narrowing the vessel lumen and reducing cerebral blood flow.¹² Additional research has also shown a correlation between the age of stroke patients and their quality of life. Older age groups often have a lower quality of life.¹⁴

This study shows that the majority of subjects, 50 out of 60, had hypertension, with 47 classified as stage II hypertension and eight as stage I hypertension. These results are consistent with previous studies in which respondents' mean blood pressure levels were placed in the hypertensive category. High blood pressure can seriously affect blood flow throughout the body, including the brain.¹⁵ Other studies have shown that patients with hypertension have a 5.69-time increased risk of having a stroke compared to those without hypertension. The occurrence of non-hemorrhagic stroke associated with hypertension is associated with cell damage in the lining of arteries, leading to the accumulation of circulating fat. This can reduce the elasticity of artery walls and impede blood circulation throughout the body, including the brain.¹³ The thickening of blood vessels and elevated blood pressure can lead to the accumulation of cholesterol and other fatty substances, resulting in the damage of arterial walls and the obstruction of cerebral arteries. Smoking and consuming foods that are rich in fat and salt worsen this condition by promoting the formation of atherosclerotic plaques. The progressive growth of atherosclerotic plaques can result in the occurrence of a stroke.¹⁶ The effects of plaque in blood vessels include narrowing of the lumen or diameter of the blood vessel. Vulnerable plaques can rupture and slough off, increasing the risk of blocking blood vessels in the brain and causing a stroke.¹⁷

The correlation analysis reveals that there is no indication of a significant correlation between blood pressure at the emergency room and mortality in patients with acute non-hemorrhagic stroke. The study results show that blood pressure ($p = 0.451$) has a significance value > 0.05 . This may be attributed to other factors influencing the mortality in acute non-hemorrhagic stroke patients. For example, the patients in this study may have received appropriate treatment to prevent a fatal outcome. Choosing a simple random sample for sample selection may also contribute to these results, as the precision and reliability of the representation are low.¹⁸ The findings of a 5-year retrospective analysis revealed that a lower Glasgow Coma Scale (GCS), decreased arterial oxygen saturation, elevated respiration rates, and increased pulse

rates at admission were correlated with the higher mortality rate of acute non-hemorrhagic stroke patients receiving treatment in hospitals. A low GCS at admission is associated with an elevated risk of mortality during the hospital period.¹⁹ The decrease in blood pressure within the initial 24 hours on admission might be used as an indicator of the probability of death in patients with acute stroke. The neurological decline and prognosis may be associated with elevated or reduced blood pressure within a 24-hour period following a stroke. Attentional observation of blood pressure within the initial 24-hour period is of the utmost importance.²⁰ Previous studies showed that the degree of stroke severity, as measured by the National Institute of Health Stroke Scale (NIHSS), correlates to a higher risk of death in individuals who have had a stroke.²¹ However, the results are conflicting, as other studies showed there is no significant correlation between initial blood pressure and final NIHSS in patients with acute thrombotic stroke.⁷

CONCLUSION

There is no significant correlation between blood pressure and mortality in patients with acute non-hemorrhagic stroke, and the correlation is very weak. Since there is no meaningful correlation between initial blood pressure and mortality in patients with acute non-hemorrhagic stroke, it is crucial to carefully monitor blood pressure over the initial 24-hour period. This is because both high blood pressure and low blood pressure may have an impact on neurological decline and potential outcomes after a stroke within the first 24 hours.

CONFLICT OF INTEREST

No conflict of interest.

REFERENCES

1. Wijaya AK. Patofisiologi Stroke Non-Hemoragik Akibat Trombus. E-J Med Udayana [Internet]. 2013 [cited 2022 Sep 30];1652–66. Available from: <https://ojs.unud.ac.id/index.php/eum/article/view/6694>
2. Mutiarasari D. Ischemic Stroke: Symptoms, Risk Factors, and Prevention. 2019;6(1):14.
3. PAPDI. Buku Ajar Ilmu Penyakit Dalam (VI). Interna Publishing; 2014.
4. Amiruddin MA, Danes VR, Lintong F. Analisa Hasil Pengukuran Tekanan Darah Antara Posisi Duduk Dan Posisi Berdiri Pada Mahasiswa Semester VII (Tujuh) TA. 2014/2015 Fakultas Kedokteran Universitas Sam Ratulangi. 2015;3:5.
5. Badriyah NJ, Amalia L, Suwarman S. Gambaran Faktor Risiko Kejadian Stroke di RSHS Bandung Periode Januari 2015–Desember 2016. J Neuroanestesi Indones [Internet]. 2018 [cited 2022 Oct 31];7(3):134–9. Available from: <http://inasnacc.org/ojs2/index.php/jni/article/view/18/pdf>
6. Rezha DM. HUBUNGAN ANTARA USIA, HIPERTENSI, KEBIASAAN MEROKOK DENGAN MORTALITAS STROKE ISKEMIK. 2019;
7. Rianawati SB, Aurora H, Nugrahanitya Y. Hubungan antara Tekanan Darah Pada Saat Masuk Stroke Unit dengan Hasil Keluaran Klinis Penderita Stroke Trombosis Akut. MNJ Malang Neurol J [Internet]. 2015 Jul 1 [cited 2022 Oct 31];1(2):51–8. Available from: <http://mnj.ub.ac.id/index.php/mnj/article/view/41>
8. Razdiq ZM, Imran Y. Hubungan antara tekanan darah dengan keparahan stroke menggunakan National Institute Health Stroke Scale. J Biomedika Dan Kesehat [Internet]. 2020 Mar 31 [cited 2023 Dec 31];3(1):15–20. Available from: <https://jbiomedkes.org/index.php/jbk/article/view/84>
9. Herdianti E, Muhartomo H, Kesoema TA. Hubungan Tekanan Darah Ketika Masuk IGD dengan Keluaran Motorik Pasien Stroke Iskemik. J Kedokt Diponegoro Diponegoro Med J [Internet]. 2018 [cited 2022 Oct 23];7(2):1604–16. Available from: <https://ejournal3.undip.ac.id/index.php/medico/article/view/21495>

10. Whelton PK, Carey RM, Aronow WS, Casey DE, Collins KJ, Dennison Himmelfarb C, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. Hypertension [Internet]. 2018 Jun [cited 2023 Dec 31];71(6). Available from: <https://www.ahajournals.org/doi/10.1161/HYP.0000000000000065>
11. Roy-O'Reilly M, McCullough LD. Age and Sex Are Critical Factors in Ischemic Stroke Pathology. Endocrinology [Internet]. 2018 Aug 1 [cited 2023 Oct 25];159(8):3120–31. Available from: <https://dx.doi.org/10.1210/en.2018-00465>
12. Nadhifah TA, Sjarqiah U. Gambaran Pasien Stroke Pada Lansia di Rumah Sakit Islam Jakarta Sukapura Tahun 2019. Muhammadiyah J Geriatr [Internet]. 2022 Aug 2 [cited 2023 Oct 25];3(1):23. Available from: <https://jurnal.umj.ac.id/index.php/MuJG/article/view/8889>
13. Azzahra V, Ronoatmodjo S. Faktor-Faktor yang Berhubungan dengan Kejadian Stroke pada Penduduk Usia >15 Tahun di Provinsi Daerah Istimewa Yogyakarta (Analisis Risesdas 2018). 2022;6(2).
14. Abdu S, Satti YC, Payung F, Soputan HA. Analisis Kualitas Hidup Pasien Pasca Stroke Berdasarkan Karakteristik. J Keperawatan Florence Nightingale [Internet]. 2022 Dec 21 [cited 2023 Oct 25];5(2):50–9. Available from: <https://ejournal.stikstellamarismks.ac.id/index.php/JKFN/article/view/107>
15. Ristonilassius, Murtiningsih, Inayah I. HUBUNGAN TEKANAN DARAH DENGAN KEJADIAN STROKE DI UNIT GAWAT DARURAT. J Keperawatan Aisyiyah [Internet]. 2022 Jul 27 [cited 2023 Oct 25];9(1):47–54. Available from: <https://journal.unisa-bandung.ac.id/index.php/jka/article/view/315>
16. Puspitasari PN. Hubungan Hipertensi Terhadap Kejadian Stroke. J Ilm Kesehat Sandi Husada [Internet]. 2020 Dec 31 [cited 2023 Oct 25];9(2):922–6. Available from: <https://akper-sandikarsa.e-journal.id/JIKSH/article/view/435>
17. Laily SR. Relationship Between Characteristic and Hypertension With Incidence of Ischemic Stroke. J Berk Epidemiol [Internet]. 2017 Apr 28 [cited 2023 Oct 25];5(1):48. Available from: <http://e-journal.unair.ac.id/index.php/JBE/article/view/3142>
18. Firmansyah D, Dede. Teknik Pengambilan Sampel Umum dalam Metodologi Penelitian: Literature Review. J Ilm Pendidik Holistik JIPH [Internet]. 2022 Aug 30 [cited 2024 Jan 1];1(2):85–114. Available from: <https://journal.formosapublisher.org/index.php/jiph/article/view/937>
19. Shah B, Bartaula B, Adhikari J, Neupane HS, Shah BP, Poudel G. Predictors of In-hospital Mortality of Acute Ischemic Stroke in Adult Population. J Neurosci Rural Pract [Internet]. 2017 Oct [cited 2023 Oct 25];08(04):591–4. Available from: <https://ruralneuropractice.com/predictors-of-in-hospital-mortality-of-acute-ischemic-stroke-in-adult-population/>
20. Niruri R, Pratika RS. Analisa Determinan Outcome Rawat Inap Pasien Stroke Iskemik Di Rumah Sakit UNS. Pros APC Annu Pharm Conf [Internet]. 2019 Nov 6 [cited 2023 Oct 25];4(1). Available from: <https://jurnal.uns.ac.id/apc/article/view/35464>
21. Sarbazi E, Sarbakhsh P, Oskooei DS, Yazdchi M, Ghaffari-Fam S, Shamshirgaran SM. Factors related to 6-month mortality after the first-ever stroke. J Educ Health Promot [Internet]. 2018 [cited 2023 Oct 25];7. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6149124/>

ORIGINALITY REPORT

18%

SIMILARITY INDEX

13%

INTERNET SOURCES

10%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1	journal.umy.ac.id Internet Source	3%
2	"Asia Pacific Stroke Conference 2018. Abstracts of the Annual Conference of the Asia Pacific Stroke Organization (APSO). Jakarta, Indonesia, September 6-9, 2018", Cerebrovascular Diseases, 2018 Publication	2%
3	Anis Ika Nur Rohmah, Marlianti Muslimah, Faqih Ruhyanudin, M. Ari Arfianto. "Factors Affecting Coping Strategies in Post-Stroke Patients: A Literature Study", KnE Medicine, 2023 Publication	1%
4	mnj.ub.ac.id Internet Source	1%
5	Submitted to Texas Woman's University Student Paper	1%
6	repository.lppm.unila.ac.id Internet Source	1%

7	Submitted to Kaplan University Student Paper	1 %
8	Submitted to University of Birmingham Student Paper	1 %
9	Submitted to Flinders University Student Paper	1 %
10	estudyassistant.com Internet Source	1 %
11	Jui-Lin Fan, Ricardo C Nogueira, Patrice Brassard, Caroline A Rickards, Matthew Page, Nathalie Nasr, Yu-Chieh Tzeng. "Integrative physiological assessment of cerebral hemodynamics and metabolism in acute ischemic stroke", Journal of Cerebral Blood Flow & Metabolism, 2021 Publication	1 %
12	Tzung-Dau Wang, Chern-En Chiang, Ting-Hsing Chao, Hao-Min Cheng et al. "2022 Guidelines of the Taiwan Society of Cardiology and the Taiwan Hypertension Society for the Management of Hypertension A Report of the Task Force of the Hypertension Committee and the Guideline Committee of the Taiwan Society of Cardiology and the Taiwan Hypertension Society", Acta Cardiologica Sinica, 2022 Publication	1 %

13	Submitted to Nightingale College - School of Nursing Student Paper	1 %
14	doi.org Internet Source	1 %
15	hdl.handle.net Internet Source	1 %
16	Jia Yu, Fude Liu, Yawen Cheng, Jianyi Wang, Wenlong Ma, Chen Chen, Peng Sun, Suhang Shang. "Burden of ischemic stroke in mainland China and Taiwan province from 1990 to 2019: with forecast for the next 11 years", International Journal for Quality in Health Care, 2023 Publication	<1 %
17	www.businessweblinks.info Internet Source	<1 %
18	perpustakaan.poltekkes-malang.ac.id Internet Source	<1 %
19	www.medicinenet.com Internet Source	<1 %
20	www.ncbi.nlm.nih.gov Internet Source	<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On