Factors that Influence Covid-19 Transmission in Patients at the Obstetrics and Gynecology Polyclinic

Ivanna Beru Brahmana¹, Elsy Maria Rosa²

¹Correspondence Author: Elsye@umy.ac.id

¹Medical Study Program, Universitas Muhammadiyah Yogyakarta, Bantul City, Indonesia

²Masters of Hospital Administration, Universitas Muhammadiyah Yogyakarta, Bantul City, Indonesia

INDEXING

Keywords:

Woman;
Suspect Covid-19;
Management
Maternal;
Obstetrician;

ABSTRACT

Stay at home is one of the health protocols for the prevention of Covid-19 transmission. The prevention is obligated for pregnant women that consult the pregnancy to the doctor for Ante Natal Care. This study aims to identify the factors that influence the transmission of covid-19 in patients who come to Obstetrics and Gynecology Polyclinic. This research method is a quantitative study with a cross-sectional survey in this study amounted to 212 respondents in the Private Practitioner Primary Service Facility. Data collection was carried out by filling out the respondent's questionnaire in May-June 2020. Data analysis using SPSS software with univariate, bivariate, and multivariate methods. The results of this study are based on the results of both univariate, bivariate, and multivariate analysis, it shows that the trips taken by the respondent within the last 1 month have an effect on the spread of covid-19 with a p-value of 0,000. Conclusion: Travel history 1 month before, influenced the Covid-19 incident with a p-value 0,000. Keywords: Women, Suspect Covid-19, Management Maternal, Obstetrician.

Kata kunci:

Perempuan;
Suspect Covid-19;
Manajemen
Maternal;
Dokter Kandungan;

Tetap di rumah adalah salah satu protokol kesehatan untuk pencegahan penularan Covid-19. Pencegahan wajib bagi ibu hamil yang mengkonsultasikan kehamilannya ke dokter untuk Ante Natal Care. Penelitian ini bertujuan untuk mengidentifikasi factor yang mempengaruhi penularan covid-19 pada pasien yang datang ke Poliklinik kebidanan dan kandungan. Metode penelitian ini adalah penelitian kuantitatif dengan survey cross sectional dalam penelitian ini berjumlah 212 responden di Fasilitas Pelayanan Pratama Praktisi Swasta. Pengumpulan data dilakukan dengan pengisian kuesioner responden pada bulan Mei-Juni 2020.. Analisis data menggunakan software SPSS dengan metode univariat, bivariat, dan multivariat. Hasil dari penelitian ini adalah berdasarkan hasil analisis baik univariate, bivariate, and multivariate menunjukkan bahwa perjalanan yang dilakukan respondent dalam kurun waktu 1 bulan terakhir berpengaruh terhadap penyebaran covid-19 dengan p-value 0,000. Kesimpulan: Riwayat perjalanan 1 bulan sebelumnya, mempengaruhi kejadian Covid-19 dengan p-value 0,000. Kata kunci: Perempuan, Suspect Covid-19, Manajemen Maternal, Dokter Kandungan.

INTRODUCTION

The World Health Organization (WHO) announced Covid-19 as a pandemic on March 11, 2020. Based on this announcement, several Health protocol rules were introduced to prevent additional casualties. The act of social distancing, which then develops into physical distancing, is applied to maintain a minimum distance of one meter when communicating with other people. Apart from that, several other protocols were also enforced, such as other measures in the form of self-isolation, staying at home, and not leaving the house if there was no very important need. This is implemented to reduce the spread of the covid-19 virus (Buekens et al., 2020). Washing your hands or using an alcohol based rub frequently and not touching your face (WHO, n.d.), and wear a mask when you leave the house or are in a crowd is one of the things that must be done during this pandemic.

The Covid-19 virus originally originated from animals, but now it is spread from humans to humans. The spread of the Corona Virus can be transmitted through several droplets, such as when coughing or sneezing, close interpersonal contact, for example touching or shaking hands, touching or rubbing the nose, mouth, and eyes before washing hands (Guan et al., 2020).

Pregnant women who need regular Ante Natal Care (ANC) checks to see an Obstetrics & Gynecology Specialist (Obsgyn) at the clinic. So that it has the potential to be infected with Covid-19. So that this problem is very important to be anticipated by doctors so that problems do not occur to mothers and babies in the delivery process. Pregnant women who are going to undergo Section Caesarea (SC) surgery need to be considered to avoid COVID-19 infection (Wong et al., 2020). The important thing to do is to prevent the spread of Covid-19 and to adhere to prevention recommendations. So, it is important to analyze complaints of pregnant or non-pregnant women, to leave the house, and come to the Obsgyn Specialist Doctor.

Pregnant women are considered vulnerable to Covid-19 infection. this could be due to physiological changes during pregnancy that have a significant impact on the immune system, respiratory system, cardiovascular function, and coagulation (Wastnedge et al., 2021). Risks that may occur in pregnant women who experience Covid-19 infection include fetal distress and preterm delivery (Chen et al., 2020). Therefore, the knowledge of health workers, especially doctors in providing clinical management in dealing with pregnant women infected with Covid-19, is an important matter that needs attention (Wang et al., 2020). Also, nurses are expected to be able to provide nursing management, especially in carrying out patient-centered care for patients infected with COVID-19 (Chiang et al., 2018).

Clinical signs and symptoms reported in most cases were fever, difficulty breathing, and X-rays showed large pneumonia infiltrate in both lungs (Kementrian Kesehatan RI, 2020). All age categories are susceptible to Covid-19 infection. Covid-19 infection is transmitted through droplets that result from coughing and sneezing by symptomatic patients, or from people without symptoms and before the onset of symptoms (Rothe et al., 2020).

The characteristics of the suspected Covid-19 case fulfill each item A and two items B, or with 3 item B as follows (Deng & Peng, 2020): 1) Epidemiological history: within two weeks before the onset of disease, have a history of travel or residence in the district with case reports, has contacted a positive patient with nucleic acid detection, have contacted a patient with fever and respiratory symptoms from a district with a reported case, and the incidence of disease in the grouping. 2) Clinical manifestations: Have fever and/or respiratory symptoms, Pneumonia with the characteristics of the images mentioned above, and in the early stages of the disease, have a normal or decreased number of leukocytes, or a decrease in the number of lymphocytes.

RESEARCH METHOD

This type of research is a quantitative study, which was conducted with a cross-sectional design in the Private Practitioner Primary Service Facility which aims to find out what complaints prompted mothers to come to Obsgyn Specialists. The data was collected by filling out a questionnaire. The clinic staff assisted in filling out the questionnaire by reading the questions posed to the subjects and then the answers were recorded by the clinic staff.

The questionnaire was submitted to subjects who agreed or were willing to participate in this research.

The subjects of this study were 212 pregnant women who came to the Specialist Doctor Obsgin in Klaten, Central Java, who were under the inclusion criteria (pregnant women) and exclusion criteria (pregnant women who had previously attended control). The research was conducted for two months, namely May and June 2020.

Data were analyzed using univariate, bivariate, and multivariate methods, to determine what factors are at risk of causing Covid-19 transmission to mothers who come to the Obsgin Specialist. The data obtained were analyzed through statistical tests used were:

1. Univariate analysis using frequency data by knowing the percentage of the characteristics of research subjects and knowing what complaints encourage mothers to come to the Obsgin Specialist Doctor.
2. Bivariate analysis by displaying the 4 variables tested which variables influence the incidence of Covid-19.
3. Multivariate analysis using multiple linear regression to determine the effect of a mother's arrival to a doctor on the incidence of Covid-19 infection.

RESULT AND DISCUSSION

The description of the research subjects is shown in Table 1 regarding the characteristics of the research subjects.

Table.1 Characteristics of Research Subjects

Characteristics	Classification	Frequency	(%)
Mother's age	< 20 th	7	3.3
	20 th - 30 th	122	57.5
	31 th - 40 th	64	30.2
	> 40 years old	19	9.0
Last education	Primary school	2	1.0
	Senior High School	17	8.0
	Diploma	84	39.6
	Bachelor	109	51.4
Mother's complaint to the clinic	ANC	135	63.7
	Post-partum	10	4.7
	Want to have children	8	3.8
	KB	13	6.1
	Post curette	2	.9
	Menstrual disorders	25	11.8
	Other gynecology	19	9
Mother's work	Professionals	66	31.1
	Non-Professional Staff	22	10.4
	Civil servants	14	6.6
	IRT	104	49.1
	Student	2	.9
	Not filled	4	1.9

Table.1 (continued) Characteristics of Research Subjects

Characteristics	Classification	Frequency	(%)
Possible Covid19	Symptomatic	30	14.2
	Asymptomatic	177	83.5
	Not filled	5	2.4
Travel History for the last 1 month	Yes	27	12.7
	Not	184	86.8
	Not filled	1	.5
	Yes	1	.5
Fever	Not	207	97.6
	Not filled	4	1.9
	Yes	3	1.4
Cough	Not	209	98.6
	Yes	0	0
Cold	Not	212	100,0
	Total	212	100,0

Source: Primary Data

Table.1 shows the characteristics of study subjects attending the clinic during a pandemic. Mass media massively reported the dangers of Covid-19 and reminded people about prevention. There are psychosocial effects and changes in individual behavior in the form of having to wear masks when leaving the house, getting used to washing hands frequently, quarantining at home, avoiding the use of public transportation, and avoiding crowds. This causes fewer mothers who come to the clinic (Yan et al., 2020). Based on the number of respondents who came to the clinic, only about 40-50% of the usual. Mothers who came to the Obsgyn Specialist Doctor during the Covid-19 pandemic were dominated by 57.5% mothers aged between 20-30 years (122/212). Mothers at that age are mothers of productive age and are still young but are psychologically mature. So they have the courage to leave the house and come to the clinic. Respondents aged under 20 years came to the clinic at least, as much as 3.3% (7/212).

Patients, clinicians, and scientists in the health sector were used to searching for information via Twitter, Facebook, and other social media platforms, even before the onset of the pandemic (Cuello-Garcia et al., 2020). The educational background of most respondents in this study, 51.4% (109/212) were undergraduate graduates. So that it is easy for them to get information through social media such as calls to leave the house only when it is necessary. The next respondent with a diploma education background was 39.6% (84/212). This shows that most mothers who dare to leave the house and come to the clinic are knowledgeable about the risks of leaving the house and consider urgent needs when leaving the house.

The effects on mental and psychosocial health are important to consider during a pandemic. In addition, there are vulnerable groups of people such as workers in the health sector, the homeless, and pregnant women (Arafat et al., 2020). This is in line with research on the reasons for mothers to come to the clinic, which is dominated by ANC needs as much as 63.7% (135/212) of pregnant women. Ante-Natal Care is a scheduled periodic prenatal examination to determine fetal development and the health of pregnant women. Thus, the need to come to an Obsgyn Specialist is a need that must be met to maintain a healthy

pregnancy until delivery. In addition, to reduce the risk of anxiety about the unwanted possibilities of fetal development.

A new social symptom in the form of panic buying often occurs during a pandemic. Anxiety causes most people to buy necessities that are not in accordance with their needs or exceed their daily needs that are usually consumed. So that in anticipating this anxiety, it is suggested that the interest to leave the house is only to fulfill daily needs and health problems (Arafat et al., 2020).

This research is in accordance with these conditions, where work as a housewife is the largest respondent with 49.1% (104/212). This is because they have more time to come to the clinic and shop for consumption needs. The next sequence was followed by professionals 31.1% (66/212), 10.4% (22/212) non-professional staff, 6.6% (14/212) civil servants, and finally students as much as 0.9% (2 / 212).

Respondents who came to the clinic were detected that 14.2% (30/212) were symptomatic of Covid-19, and 83.5% (177/212) had no symptoms, and there were 5 data that were not filled in. As many as 12.7% (27/212) of respondents had a history of traveling in the last month, and 86.8% (184/212) had not traveled, but it was found that 0.5% (1/212) did not fill in data. The results showed that 0.5% (1/212) had a fever, 97.6% (207/212) and 1.9% (4/212) had no fever. Complaints of cough were experienced by 1.4% (3/212) and 98.6% (209/212) of coughs, while 100% of respondents did not complain of colds (212/212). Respondents are at risk for symptoms of Covid-19 or not, a history of travel before coming to the clinic, especially from the red zone, complaints of fever, cough, colds need to be disclosed in this study because it protects health workers working in clinics, who interact directly with patients (Mhango et al., 2020).

In line with the research of Ogolodom MP. et al. 68% (204/300) of respondents know that the complaints that arise in Covid-19 sufferers are fever more than 380C, cough, shortness of breath, weakness, nasal congestion, nausea, diarrhea, and sneezing (Mp et al., 2020). This is following Remuzzi's research, cited by Fadli et al., That the main symptoms of Covid-19 are complaints of fever, weakness, cough, and diarrhea (Fadli et al., 2020).

Table.2 Results Of Bivariate Analysis

Variable	P-Value Possible Covid-19	Result
Mother's job	0.006	Significant
Mother's age	0.333	Not significant
Last education	0.025	Significant
Travel History for the last 1 month	0.000	Significant

The results of the bivariate analysis showed that the variables of employment, education, and travel history in the last month had a significant value for the risk of Covid-19, with p values, respectively: 0.006, 0.025, and 0.000. Maternal age is not a significant risk for Covid-19 with a p-value of 0.333. A good recording system as a data system needed for the latest daily reports is needed during a sudden pandemic like Covid-19. This is in line with this study, where when respondents came to the clinic, their consent was asked to fill in data about their identity and travel data expressed their enthusiasm and consent (Jafar & Fletcher, 2020).

The results of the multivariate analysis using the Multiple Linear Regression Test regarding the likelihood of being infected with Covid-19 in mothers who came to the clinic were influenced by variables: age (X1), education (X2), occupation (X3), and travel history in the last one month (X4), obtained the following results:

R Square Value (R²)

Table.3 The Value of R Square (R²) Effect of Age, Education, Occupation, and Travel History on The Likelihood of Being Infected with Covid-19

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	.918 ^a	.843	.840	.140

Table.3 states that R Square is 0.843 (84.3%). This shows that the possibility of Covid-19 occurring in mothers who come to see an Obsgyn Specialist is influenced by variables of age, education, occupation, and travel history in the past month. Other things apart from that of 0.157 (15.7%) are influenced by other factors, outside the four variables, studied. The factors of age, education, and work show an influence on the possibility of Covid-19 in this study in line with the research of Chatterjee R., et al., Which explains about access to data about Covid-19 using a mobile phone, the majority of respondents are aged between 10-39 years, followed by those aged 40-49 years (Chatterjee et al., 2020). In that study, 853/2216 respondents were 10-39 years old. In this study, most respondents who came to the clinic were 20-30 years old.

This is also in line with the research of Barua Z., et al., where the respondents were mostly between the ages of 18-40 years as many as 365/483, compared to those who were over 40 years as many as 118/483 respondents (Barua et al., 2020).

F test (Anova)

The results of the analysis of the effect of four variables (age (X1), education (X2), occupation (X3), and travel history (X4)) on the possibility of being infected with Covid-19 together can be seen in table 4.

Table.4 Effect of Variables (Age (X1), Education (X2), Occupation (X3), and Travel History (X4)) on the Possibility of Being Infected with Covid-19

ANOVA^a

Model	Sum of Squares	df	Mean Square	F	Sig.
1 Regression	20.962	4	5.240	266.383	.000 ^b
Residual	3.895	198	.020		
Total	24.857	202			

The Anova test table shows that the four variables (age (X1), education (X2), occupation (X3), and travel history (X4)) on the likelihood of being infected with Covid-19 have a significant effect on the likelihood of Covid-19 occurring, with a p value 0,000.

T-Test**Table.5 T-Test Values on 4 Variables**

Model	Unstandardized Coefficients		Standardized Coefficients	t	Sig.
	B	Std. Error	Beta		
1 (Constant)	-.006	.083		-.075	.940
Age	-.002	.014	-.004	-.154	.878
Profession	-.002	.008	-.010	-.320	.749
Education	.005	.006	.021	.706	.481
Travel History of the Last 1 Month	.984	.031	.924	31.635	.000

Travel history for the past month has a very significant p-value of 0.000, compared to the other three variables which are not significant. In this study, travel history for the past month has significant risk factors being infected Covid-19. A study reported that staying at home would prevent infection with Covid-19 around 96 (32%) (Mp et al., 2020).

The regression equation to explain the effect of age, education, occupation, and travel history for the last month on the variable likelihood of being infected with Covid-19 is as follows:

$$Y = -0,004 X1 - 0,010 X2 + 0,021 X3 + 0,924 X4 + e$$

X1 = Age.

X2 = Work.

X3 = Education.

X4 = Travel History of the Last Month.

CONCLUSION

Pregnant women have physiological changes during pregnancy that have a significant impact on the immune system, respiratory system, and cardiovascular function, which makes pregnant women vulnerable to Covid-19 infection. Government measures to prevent the spread of the Covid-19 virus, such as limiting physical distance and travel. Respond to this preventive measure is quite good in the community, especially pregnant women, but their willingness to implement it is still low. The limitation in this study is that pregnant women as housewives force them to travel outside the home to meet household needs.

ACKNOWLEDGMENT

The authors would like to thank UMY for providing funds for the implementation of this research.

REFERENCE

- Arafat, S. M. Y., Kar, S. K., Menon, V., Kaliamoorthy, C., Mukherjee, S., Alradie-Mohamed, A., Sharma, P., Marthoenis, M., & Kabir, R. (2020). Panic buying: An insight from the content analysis of media reports during COVID-19 pandemic. *Neurology, Psychiatry, and Brain Research*, 37, 100–103. <https://doi.org/10.1016/j.npbr.2020.07.002>
- Barua, Z., Barua, S., Aktar, S., Kabir, N., & Li, M. (2020). Effects of misinformation on COVID-19 individual responses and recommendations for resilience of disastrous

- consequences of misinformation. *Progress in Disaster Science*, 8, 100119. <https://doi.org/10.1016/j.pdisas.2020.100119>
- Buekens, P., Alger, J., Bréart, G., Cafferata, M. L., Harville, E., & Tomasso, G. (2020). A call for action for COVID-19 surveillance and research during pregnancy. *The Lancet Global Health*, 8(7), e877–e878. [https://doi.org/10.1016/S2214-109X\(20\)30206-0](https://doi.org/10.1016/S2214-109X(20)30206-0)
- Chatterjee, R., Bajwa, S., Dwivedi, D., Kanji, R., Ahammed, M., & Shaw, R. (2020). COVID-19 Risk Assessment Tool: Dual application of risk communication and risk governance. *Progress in Disaster Science*, 7, 100109. <https://doi.org/10.1016/j.pdisas.2020.100109>
- Chen, R., Zhang, Y., Huang, L., Cheng, B.-H., Xia, Z.-Y., & Meng, Q.-T. (2020). Safety and efficacy of different anesthetic regimens for parturients with COVID-19 undergoing Cesarean delivery: A case series of 17 patients. *Canadian Journal of Anaesthesia = Journal Canadien D'anesthésie*, 67(6), 655–663. <https://doi.org/10.1007/s12630-020-01630-7>
- Chiang, C.-Y., Choi, K.-C., Ho, K.-M., & Yu, S.-F. (2018). Effectiveness of nurse-led patient-centered care behavioral risk modification on secondary prevention of coronary heart disease: A systematic review. *International Journal of Nursing Studies*, 84, 28–39. <https://doi.org/10.1016/j.ijnurstu.2018.04.012>
- Cuello-Garcia, C., Pérez-Gaxiola, G., & van Amelsvoort, L. (2020). Social media can have an impact on how we manage and investigate the COVID-19 pandemic. *Journal of Clinical Epidemiology*, 127, 198–201. <https://doi.org/10.1016/j.jclinepi.2020.06.028>
- Deng, S.-Q., & Peng, H.-J. (2020). Characteristics of and Public Health Responses to the Coronavirus Disease 2019 Outbreak in China. *Journal of Clinical Medicine*, 9(2). <https://doi.org/10.3390/jcm9020575>
- Fadli, F., Ahmad, A. S., Safruddin, S., Sumbara, S., & Baharuddin, R. (2020). Anxiety of Health Workers in the Prevention and Management of Covid-19 in Sidrap Regency. *Unnes Journal of Public Health*, 9(2), 91–97. <https://doi.org/10.15294/ujph.v9i2.38436>
- Guan, W., Liang, W., Zhao, Y., Liang, H., Chen, Z., Li, Y., Liu, X., Chen, R., Tang, C., Wang, T., Ou, C., Li, L., Chen, P., Sang, L., Wang, W., Li, J., Li, C., Ou, L., Cheng, B., ... He, J. (2020). Comorbidity and its impact on 1590 patients with COVID-19 in China: A nationwide analysis. *The European Respiratory Journal*, 55(5). <https://doi.org/10.1183/13993003.00547-2020>
- Jafar, A. J. N., & Fletcher, R. J. (2020). Using field hospital simulation to demonstrate changes in completion rates of a UK emergency medical team (EMT) medical record. *Progress in Disaster Science*, 6, 100072. <https://doi.org/10.1016/j.pdisas.2020.100072>
- Kementrian Kesehatan RI. (2020). *Pedoman Pencegahan dan Pengendalian COVID-19*.
- Mhango, M., Dzobo, M., Chitungo, I., & Dzinamarira, T. (2020). COVID-19 Risk Factors Among Health Workers: A Rapid Review. *Safety and Health at Work*, 11(3), 262–265. <https://doi.org/10.1016/j.shaw.2020.06.001>

- Mp, O., An, M., N, A., Of, E., No, E., I, G., Dc, U., Gi, A., & Cm, E. (2020). Knowledge, Attitudes and Fears of HealthCare Workers towards the Corona Virus Disease (COVID-19) Pandemic in South-South, Nigeria. *Health Science Journal*, 14(0). <https://doi.org/10.36648/1791-809X.S1.002>
- Rothe, C., Schunk, M., Sothmann, P., Bretzel, G., Froeschl, G., Wallrauch, C., Zimmer, T., Thiel, V., Janke, C., Guggemos, W., Seilmaier, M., Drosten, C., Vollmar, P., Zwirgmaier, K., Zange, S., Wölfel, R., & Hoelscher, M. (2020). Transmission of 2019-nCoV Infection from an Asymptomatic Contact in Germany. *New England Journal of Medicine*, 382(10), 970–971. <https://doi.org/10.1056/NEJMc2001468>
- Wang, S., Zhou, X., Lin, X., Liu, Y., Wu, J., Sharifu, L. M., Hu, X., Rong, Z., Liu, W., Luo, X., Chen, Z., Zeng, W., Chen, S., Ma, D., Chen, L., & Feng, L. (2020). Experience of Clinical Management for Pregnant Women and Newborns with Novel Coronavirus Pneumonia in Tongji Hospital, China. *Current Medical Science*, 1–5. <https://doi.org/10.1007/s11596-020-2174-4>
- Wastnedge, E. A. N., Reynolds, R. M., van Boeckel, S. R., Stock, S. J., Denison, F. C., Maybin, J. A., & Critchley, H. O. D. (2021). Pregnancy and COVID-19. *Physiological Reviews*, 101(1), 303–318. <https://doi.org/10.1152/physrev.00024.2020>
- WHO. (n.d.). Coronavirus. Retrieved December 30, 2020, from <https://www.who.int/westernpacific/health-topics/coronavirus>
- Wong, J., Goh, Q. Y., Tan, Z., Lie, S. A., Tay, Y. C., Ng, S. Y., & Soh, C. R. (2020). Preparing for a COVID-19 pandemic: A review of operating room outbreak response measures in a large tertiary hospital in Singapore. *Canadian Journal of Anaesthesia = Journal Canadien D'anesthesie*, 67(6), 732–745. <https://doi.org/10.1007/s12630-020-01620-9>
- Yan, Q., Tang, Y., Yan, D., Wang, J., Yang, L., Yang, X., & Tang, S. (2020). Impact of media reports on the early spread of COVID-19 epidemic. *Journal of Theoretical Biology*, 502, 110385. <https://doi.org/10.1016/j.jtbi.2020.110385>