Histopathological Evaluation of Soybean (*Glycine max* (L.) Merr.) Strains Resistance to *Sclerotium rolfsii* Disease

DOI: [10.18196/pt.v10i1.8907](https://doi.org/10.18196/pt.v10i1.8907)

Siti Samiyarsih^{1*}, Chindy Ayu Erliana¹, Juni Safitri Muljowati², Nur Fitrianto³

¹Department of Botany, Faculty of Biology, Universitas Jenderal Soedirman

²Department of Microbiology, Faculty of Biology, Universitas Jenderal Soedirman, Jalan Dr. Suparno No. 63 Purwokerto 53122, Banyumas, Central Java, Indonesia

³Research and Technology Center for Application of Isotope and Radiation, National Research and Innovation Agency of Indonesia (BRIN); Jl. Lebak Bulus Raya No. 49 Jakarta 12440, Indonesia.

*Corresponding author, email: siti.samiyarsih@unsoed.ac.id

ABSTRACT

Sclerotinia infection of stem and leaf of soybean *Glycine max* (L.) Merr. caused by the fungal pathogen of *Sclerotium rolfsii* has recently become more important in the Indonesian soybean production area. This study aimed to evaluate the level of resistance and intensity of infection by *S. rolfsii* in four soybean strains. The research was arranged in a factorial completely randomized design. The observed variables include the anatomy characteristics of leaves and stems of soybean and disease intensity caused by *S. rolfsii*. The data were analyzed quantitatively with the Analysis of Variance (ANOVA) at 95% and 99% confidence level, followed by the Least Significant Difference Test (Fisher's LSD) at the level of 5%. Soybean leaves and stem anatomy inoculated by *S. rolfsii* showed a decrease in the stomatal density, epidermis thickness, and mesophyll thickness as well as a damaged cuticle, damaged stem epidermis, and swollen stem cortex. Four strains inoculated by *S. rolfsii* showed a higher disease intensity of 40%-80% compared to the resistant cultivar ('Dering') and susceptible cultivar ('Wilis'), showing disease intensity of 20% and 40%, respectively.

Keywords: *Glycine max*, Histopathology, Resistance, *Sclerotium rolfsii*

ABSTRAK

Infeksi sclerotinia pada daun dan batang kedelai *Glycine max* (L.) Merr. yang disebabkan oleh jamur patogen *Sclerotium rolfsii* menjadi semakin penting di area produksi kedelai Indonesia. Penelitian ini bertujuan mengevaluasi tingkat resistensi dan tingkat intensitas infeksi oleh *S. rolfsii* terhadap empat galur kedelai. Metode yang digunakan dalam penelitian ini adalah metode eksperimen dengan pola rancangan acak lengkap pola faktorial. Parameter yang diamati meliputi karakteristik anatomi pada daun dan batang kedelai, dan intensitas penyakit yang disebabkan oleh *S. rolfsii*. Data dianalisis dengan Analisis Varians (ANOVA) dengan tingkat kepercayaan 95% dan 99%, analisis data dilanjutkan dengan Uji Beda Nyata Terkecil (LSD) 5%. Karakteristik histopatologi daun dan batang yang diinokulasi oleh *S. rolfsii* menunjukkan adanya penurunan kerapatan stomata, ketebalan epidermis dan ketebalan mesofil, sekaligus menyebabkan kerusakan pada lapisan kutikula, epidermis batang, dan pembengkakan pada korteks batang. Empat galur yang diinokulasi oleh *S. rolfsii* memiliki intensitas penyakit yang lebih tinggi yaitu 40% -80% dibandingkan dengan kultivar 'Dering' sebagai kelompok tahan sebesar 20% dan kelompok rentan pada kultivar 'Wilis' sebesar 40%.

Kata kunci: *Glycine max*, Histopatologi, Resistensi, *Sclerotium rolfsii*

INTRODUCTION

Soybean (*Glycine max* L.) is an important protein source plant in Indonesia. The increase of soybean production is in line with the increase in the number of population and industrial developments using soybean as raw materials. According to Wahyu (2013), soybean is one of the widely cultivated legume commodities in Indonesia. Due to the high consumption rate, efforts are needed to increase soybean production through superior cultivars. However, efforts to increase soybean

production can not be separated from various obstacles, including pest and disease attacks. One of the acute diseases is stem rot disease caused by the *Sclerotium rolfsii*.

Histopathology based on leaves anatomy characters can be used as instructions to the structural resistance of plants against the pathogen (Samiyarsih et al., 2018). *S. rolfsii* is a fungus that can cause several diseases in plants, such as stem rot. A wilting disease caused by *S. rolfsii* is a common

Article History
Received: 28 May 2020
Accepted: 31 Jan 2022

Planta Tropika: Jurnal Agrosains (Journal of Agro Science) is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

disease in soybean plants. This disease is often also referred to as stem rot disease or sclerotium rot because it causes root rot symptoms. *S. rolfsii* can also attack leaves, stems, and pods in soybean plants if conditions are very humid. Efforts to get high-yielding soybeans can pursue through plant breeding activities. Plant breeding is expected to improve and increase plants' genetic potential so that superior results are obtained with suitable characteristics, one of which is by selecting new strains (Wardoyo, 2009). Pure strain selection is made by choosing the best plants. Selected plants are individually harvested separately for planting material the following season. Cultivar differences in plants will provide genetically different responses (Sasongko et al., 2019).

The histopathology based on plants' anatomical structure plays an essential role in the relationship with pathogenic infections in tissues. The disease's effects cannot adequately be understood without understanding the typical structure of the affected tissue. Besides, the impact of disease or parasites and even susceptibility to disease can be identified by structural changes in the host structure's characteristics. The primary response of plants affected by fungal infections is structural defenses, such as cell wall thickness. The disease can prevent the penetration of pathogens into the host cell. Besides, pathogenic infections can cause the development of plant vascular tissue structures to be disrupted (Impullitti et al., 2014).

In this research, pure soybean strains were used to determine each different soybean strain's anatomical characteristics then compared to the cultivars. This study aimed to determine the differences in the anatomical character of leaves and stems in soybean strains that are resistant and susceptible to stem rot disease after inoculation with *S. rolfsii* and the level of intensity of attacks of each soybean strain used after *S. rolfsii* fungal infection.

MATERIALS AND METHODS

The research was conducted in June-September 2019 at the greenhouse, the Plant Structure and Development Laboratory, and the Mycology and Phytopathology Laboratory, Faculty of Biology, Jenderal Soedirman University, Puwokerto. This research was arranged in a completely randomized factorial design. The first level was four soybean strains (Strain no. 71-7, 39-6, 32-6, and 16-4) and two types of soybean cultivars ('Dering' and 'Wilis'). Soybean strains and cultivars are the collections of the Faculty of Agriculture, Jenderal Soedirman University. The second level was fungal inoculation treatment. Tests without inoculation and with fungal inoculation were carried out with five replications.

The isolates of *S. rolfsii* cultures were propagated in a potato dextrose agar (PDA) medium and incubated at room temperature for 5×24 h (Astiko, 2009). Propagation of fungal inoculums was done using bran media as much as $\frac{3}{4}$ the volume of the bottle that had been sterilized first in the autoclave then inoculated with three plugs of rejuvenated fungal mycelium on PDA medium (Nugroho, 2008). The inoculation of *S. rolfsii* was carried out on the 14-day-old plant in the polybag. The inoculation of *S. rolfsii* was carried out by giving an inoculum at a depth of ± 1 cm between the plant's roots and on the soil surface.

Anatomical characteristics including cuticle, epidermis, mesophyll, stomata size, stomata, and trichomes density per mm² area of the epidermis of leaves were observed using embedding methods of Khoiroh et al. (2014) and Samiyarsih et al. (2020a) with a slight modification. The 5th leaf from the shoot bud was taken and cut into one-cm pieces. It was subjected to fixation in FAA solution (FAA: 10% formalin, 5% acetic acid, 50% ethyl alcohol, and 35% distilled water) for 24 h. Staining was done using safranin (1%) in 70% alcohol. Observa-

tion of anatomical characteristics was performed using a binocular microscope Olympus CH-20 (Damayanti, 2007). The incubation period was from the first day after the inoculation of pathogenic fungi until the disease's symptoms appeared. Observation of disease intensity was carried out to determine the level of resistance of soybean plants to stem rot disease. Consideration of the severity of the disease was carried out 15 days after planting. The percentage of disease intensity could be calculated using the formula of $I = N / n \times 100\%$, in which I is the severity of the disease, n is the number of plants showing symptoms, and N is the number of plants observed. Disease intensity calculation results were then categorized based on the assessment of the level of resistance (Disease intensity (%) resistance: 0-20=high; >20-30=medium, and > 30=low). All data were analyzed by ANOVA followed by Least Significant Difference (LSD) test at 0.05 (5%).

RESULTS AND DISCUSSION

Leaves Histopathological Characteristics of Soybean Strains Affected by *S. rolfsii*

The histopathology of soybean leaves that were attacked by *S. rolfsii* causing stem rot disease in the 'Dering' and 'Wilis' cultivars and four soybean strains with strain numbers 71-7; 39-6; 32-6; and 16-4 showed the similar leaf anatomical structure (Table 1; Figure 1). The plant inoculated by *S. rolfsii* was damaged in the leaf epidermis and leaf mesophyll tissue. Meanwhile, a reduced palisade density characterized damage to the mesophyll tissue, and the space between cells contained in spongy tissue or sponges had also become more extensive. The strain no. 39-6 showed the least tissue damage compared to other strains and cultivars.

The highest thickness of the epidermis was observed in strain number 39-6, with a thickness of the adaxial and abaxial epidermis of 10.1 and 9 μm , respectively. Meanwhile, the highest meso-

Table 1. Histopathological characteristics of leaves and stem caused by *S. rolfsii* disease

No	Cultivar/Strain	Adaxial epidermis thickness	Abaxial epidermis thickness	Mesophyll thickness	The adaxial density of stomata	The abaxial density of stomata	The abaxial density of trichomes	The abaxial density of trichomes	Stem diameter
1	Dering	9.05 c	9.30 abc	58.65 d	5.40 a	8.40 bc	5.40 a	8.40 bc	2597.4 b
2	Wilis	8.50 c	8.50 cd	57.00 d	4.92 ab	7.72 bc	4.92 ab	7.72 bc	2524.8 b
3	Strain no. 71-7	8.60 c	8.25 d	61.95 d	4.96 ab	8.36 bc	4.96 ab	8.36 bc	3089.8 a
4	Strain no. 39-6	10.50 a	10.10 a	95.90 b	5.16 ab	10.00 a	5.16 ab	10.00 a	2652.0 b
5	Strain no. 32-6	10.10 ab	9.50 ab	73.60 c	4.54 b	8.96 ab	4.54 b	8.96 ab	2483.0 b
6	Strain no. 16-4	9.25 bc	8.85 bcd	104.30 a	3.86 c	7.30 c	3.86 c	7.30 c	2496.8 b

Remarks: Values followed by the same letters are not significantly different according to LSD at 5%.

phyll thickness was found in strain number 16-4 (103.7 μm), and the lowest mesophyll thickness was in the 'Wilis' cultivar (48.6 μm). The highest number of adaxial epidermal stomata was in strain number 39-6 (5.16/ mm^2), and the lowest was in strain number 16-4 (3.24/ mm^2). The same results were obtained in the number of lower epidermal stomata. The highest and lowest number of lower epidermal stomata was observed in strain number 39-6 (10.16/ mm^2) and strain number 16-4 (6.44/

mm^2), respectively. There was significant difference in the number of adaxial and abaxial epidermal stomata (Figure 1). Kouwenberg et al. (2004) noted that morphogenesis changes caused variations in stomatal density between plants of various dicotyledonous plant species. Environmental adaptation factors can also influence the calculation of the number of stomata. Juwarno et al. (2017) reported that the adaxial and abaxial stomata density was not significantly different between soybean cultivars.

Figure 1. Leaves histopathology of soybean strains for resistance to *S. rolfisii* disease. Notes: A1-F1 (uninfected with *S. rolfisii*); A2-F2 (Infected with *S. rolfisii*); (A) Cultivar 'Dering'; (B) Cultivar 'Willis'; (C) Strain no 71-7; (D) Strain no. 39-6; (E) Strain no. 32-6; (F) Strain no. 16-4; (e.a) adaxial epidermis ; (p) palisade; (s) sponges; (e.b) abaxial epidermis.

The density of trichomes indicated less intensity of leaf damage. Based on different test tables, strain no. 39-6 showed similar average number of stomata to the resistant cultivar 'Dering'. Meanwhile, strain no. 32-6 and strain no. 16-4 had lower number of stomata compared to the susceptible cultivar 'Willis'. The average number of trichomes in the upper epidermis was less than that of the lower epidermis. Wijaya (2016) states that the difference in the number and length of trichomes on the adaxial and abaxial surfaces of leaves is influenced by plant genetic factors to prevent pests and diseases that usually attack through the underside of the leaves. Arifin (2013) adds that the number of trichomes in healthy soybean plants is higher more than in sick soybean plants. Pradana et al. (2017) reported that the density of stomata-trichomes was the same as the plant disease intensity. On the other hand, Samiyarsih et al. (2020b) mention that soybean

cultivars that have thicker cuticle and epidermis, high trichomes and low stomatal density, and low stomatal conductance have better anatomical resistance to leaf rust disease.

The lowest thickness of the upper epidermis was found in the 'Willis' cultivar (5.8 μm), but the lowest epidermis thickness was observed in the strain 71-7 (6 μm). Decreased epidermal thickness is thought to occur as a result of changes in cell permeability in response to pathogens. Sastrahidayat (1989) summarized that reduced cell permeability was the beginning of changes in diseased tissue. The cells in the tissue that is attacked and damaged often undergo plasmolysis. Besides, the decrease in plants' epidermal thickness inoculated with *S. rolfisii* can also be caused by chemicals during preparations. Diseased plant tissue is more easily damaged when given treatment using chemicals. According to Samiyarsih et al. (2018), disease-resistant plants

Figure 2. Stems histopathology of soybean strains for resistance to *S. rolf sii* disease. Notes: A1-F1 (uninfected with *S. rolf sii*); A2-F2 (Infected with *S. rolf sii*); (A) Cultivar 'Dering'; (B) Cultivar 'Wilis'; (C) Strain no 71-7; (D) Strain no. 39-6; (E) Strain no. 32-6; (F) Strain no. 16-4; (e.a) adaxial epidermis ; (p) palisade ; (s) sponges; (e.b) abaxial epidermis.

tend to have thick epidermis, playing an essential role in inhibiting pathogens penetration into host cells.

Stem Diameter of Soybean Strains Affected by *S. rolf sii*

Strain number 71-7 were significantly different from the cultivar 'Dering,' cultivar 'Wilis,' strain number 39-6, strain number 32-6, and strain no. 16-4. Other than strain numbers 71-7, the other strains and cultivars had uniform stem size (Figure 2). Stems infected with *S. rolf sii* showed a more brownish color. The color is due to the attacks of *S. rolf sii* to the stem's base, causing the bottom

of the stem to be swollen before finally decaying. Phenol accumulation also occurs, causing the stem to turn brown. Changes in metabolism in diseased plants accompany an increase in respiration after infection because the enzymes associated with respiration increase. Tang et al. (2015) reported that *S. rolf sii* produced a variety of extracellular enzymes, including pectin methylesterase, cutaneous, phosphatides, arabanase, gateringase, mannanase, xylanase, oxalic acid, and polygalacturonase, which are thought to cause tissue death along with mycelial growth during the infectious process.

The stem diameter of the cultivar 'Dering' inoculated with *S. rolf sii* had a larger size than that

Figure 3. Percentage of disease intensity of soybean strains against *S. rolfsii* (in three replication).

of the cultivar ‘Dering’ without inoculation. This is related to the swollen stems of plants after the inoculation of *S. rolfsii*. It was characterized by an enlarged size of the cortex of stems infected with *S. rolfsii* due to disruption of nutrient absorption by xylem and phloem compared to stems uninfected with *S. rolfsii*. The cortex’s internal part is a system of stem vessels consisting of the phloem on the outside and xylem. Another factor that is thought to have an influence is the swollen cell wall, which increases diameter. Direct penetration occurs in the epidermis cell wall by a pathogen, and sometimes the outer cell wall will be swollen, thereby inhibiting pathogen penetration. The results of the analysis of the variety of soybean stem diameters showed very significant values. Pranita et al. (2010) investigated that in stems experiencing secondary growth, the epidermal layer is replaced by a cork layer formed from cork cambium. The cork layer in plants helps increase the protective power of the stem and reduce water evaporation.

Pathogenicity of *S. rolfsii* to soybean strains and cultivars

The pathogenicity test results on soybean revealed that *S.rolfsii* fungi were capable of infecting test plants, including the ‘Dering’ cultivar, ‘Wilis’ cultivar, strain no. 71-7, strain no. 39-6, strain no.

32-6, and strain no. 16-4. ‘Dering’ cultivar showed the lowest disease intensity value of 20%. ‘Wilis’ cultivar and line no. 39-6 had the same disease intensity value, equal to 40%, and strain no. 71-7 showed a disease intensity of 60%. Meanwhile, the highest disease intensity was found in strain-no-32-6, which is 80% (Figure 3). The disease intensity of ‘Dering’ cultivar showed a high level of disease resistance. Meanwhile, the ‘Wilis,’ *S. rolfsii* cultivar, strain no. 71-7, strain no. 39-6, line 32-6, and line no. 16-4 are categorized as low resistance.

In soybean plants, the symptom is the start of the withering of soybean plants, accompanied by the stems’ base that begins to rot. The incubation period for *S. rolfsii* pathogens in test plants ranged from three to nine days. Environmental factors provide a considerable influence for pathogens to infect soybean plants. Environmental conditions due to routine watering in the morning and evening cause the soil around the stems to become more humid. This is undoubtedly beneficial for the breeding of *S. rolfsii* spores. Sumartini (2011) reported that the *S. rolfsii* would be more infective at high humidity, causing high intensity and extent of the attack. Conversely, low moisture would stimulate *S. rolfsii* to form sclerotia.

The novelty of selecting soybean germplasm

against biotrophic fungal disease is essential and effective in order to increase crop productivity (Samiyarsih et al., 2020b). Overall, the level of resistance in the four strains observed is relatively low. This is due to the high intensity of disease in the four strains compared to cultivars that have been released, which is above 30%. The higher the intensity of the disease, the lower the resistance to pathogens. Astiko et al. (2009) stated that there were differences in resilience plants among soybean varieties in suppressing the development of rot disease stem base. Each strain has different resistance characteristics to *S. rolfii* attacks due to the different resistance genes controllers to fight pathogens in each variety.

CONCLUSION

The difference in disease intensity of soybean strains and cultivars tested is greatly influenced by plant resistance. Histopathological evaluation of soybean leaves inoculated by *S. rolfii* showed decreased leaf epidermis thickness and leaf mesophyll thickness, as well as damage to the cuticles, stem epidermis, and swollen cortex of the stem. All Strains inoculated with *S. rolfii* showed a higher disease intensity of 40% -80% compared to the resistant cultivar 'Dering' (20%), and the susceptible cultivar 'Willis' (40%). This method is helpful in differentiating reactions of soybean strain or cultivars to *S. rolfii*.

REFERENCES

- Arifin, A. S. (2013). Kajian Morfologi Anatomi dan Agronomi antara Kedelai Sehat dengan Kedelai Terserang Cowpea Mild Mottle Virus serta Pemanfaatannya sebagai Bahan Ajar Sekolah Menengah Kejuruan. *Jurnal Pendidikan Sains*, 1(2), pp. 115-125.
- Astiko, W., Muthahanas, I., & Fitrianti, Y. (2009). Uji Ketahanan Beberapa Kultivar Kacang Tanah Lokal Bima Terhadap Penyakit *Sclerotium rolfii* Sacc. *Crop Agro*, 5(1).
- Damayanti, F. (2007). Analisis Jumlah Kromosom dan Anatomi Stomata pada Beberapa Plasma Nutfah Pisang (Musasp.) Asal Kalimantan Timur. *Journal of Bioscientiae*, 4(2), pp. 53 - 61.
- Impullitti, A. E & Malvick, D. K. (2014). Anatomical Response and Infection of Soybean during Latent and Pathogenic Infection by Type A and B of *Phialophora gregata*. *PLOS ONE*, 9(5), pp.1-11.
- Juwarno, J., & Samiyarsih, S. (2017). Anatomical and Molecular Responses of Soybean (*Glycine Max* (L.) Merr.) Due to Salinity Stresses. *Molekul*, 12(1): 45-52.
- Khoiroh Y, Harijati N & Retno M. (2014). Pertumbuhan Serta Hubungan Kerapatan Stomata dan Berat Umbi pada *Amorphophallus muelleri* Blume. & *Amorphophallus variabilis* Blume. *Jurnal Biotropika*, 2 (5): 249-253.
- Kouwenberg, L. L. R., Kurschner, W. M., & Visscher, H. (2004). Changes in Stomatal Frequency and Size During Elongation of *Tsuga heterophylla* Needles. *Annals of Botany*, 94, pp. 561-569.
- Nugroho, B. A. (2008). Cara Membuat Media Tumbuh dalam Pengembangan Massal APH Golongan Jamur. Surabaya: POPT BBP2TP.
- Pradana, A. W., Samiyarsih, S., & Muljowati, J. S. (2017). Korelasi karakter anatomi daun ubi jalar (*Ipomoea batatas* L.) kultivar tahan dan tidak tahan terhadap intensitas penyakit kudis daun. *Scripta Biologica*, 4(1), 21-29.
- Pranita, R., Y. R. Fitri, T. Asneti, Juwilda, E. F. Zeba. (2010). Epidermis Pada Tumbuhan Makalah. Universitas Sriwijaya: Inderalayah.
- Samiyarsih S, Fitrianto N, Azizah E, Herawati W, Rochmatino. (2020a). Anatomical profile and genetic variability of sweet potato (*Ipomoea batatas*) cultivars in Banyumas, Central Java, based on RAPD markers. *Biodiversitas* 21(4): 1755-1766.
- Samiyarsih, S., Pratiwi, A. Y. P., Muljowati, J. S., & Fitrianto, N. (2020b). Selection of Soybean (*Glycine max*) Germplasm Against Biotrophic Fungi Disease Based on Anatomical Resistance. *Biosaintifika: Journal of Biology & Biology Education*, 12(3): 311-318.
- Sasongko ND, Samiyarsih S, Juwarno. (2019). Genetic variation among the scabies-infested sweet potato cultivars. *Intl J CurrRes* 11 (7): 5115-5120.
- Sumartini. (2011). Penyakit Tular Tanah (*Sclerotium rolfii* dan *Rhizoctonia solani*) pada Tanaman Kacang-Kacangan dan Umbi-Umbian Serta Cara Pengendaliannya. *Jurnal Litbang Pertanian*, 31(1), pp. 1-8.
- Tang, Wei., Kuang, J., & Qiang, S. (2015). The Pathogenicity of *Sclerotium rolfii* on *Cyperus difformis* and its Potential Host Specificity among the Genus *Cyperus*. *Journal Plant Pathology & Microbiology*, 8(3): 1-6.
- Wahyu, E. R., Purwani, K. I., & Nurhatika, S. (2013). Pengaruh *Glomus fasciculatum* Pada Pertumbuhan Vegetatif Kedelai yang Terinfeksi *Sclerotium rolfii*. *Jurnal Sains dan Seni POMITS*, 2(2): 1-5.
- Wardoyo, S. D. W. (2009). Uji Daya Hasil Lanjutan Galur-Galur Harapan Kedelai (*Glycine max* (L.) Merr.) Berdaya Hasil Tinggi. Skripsi. Fakultas Pertanian IPB. Bogor.
- Wijaya, I., Zubaidah, S., & Kuswantoro, H. (2016). Karakter Anatomi Galur-Galur Harapan Kedelai (*Glycine max* L. Merrill) Tahan Cowpea Mild Mottle Virus (CpMMV). *Jurnal Pendidikan Biologi Bioedukasi*, 7(1): 16-25.